

NRD Trustees Handouts

- [Jan 2013- Hudson River Status Report Update](#)
- [August 2013- Mink Modification Report](#)
- [Sept 2013- Hudson River Fact Sheet Restoration Project Proposals List](#)
- [Sept 2013- Mussel Study Fact Sheet](#)

PCB CONTAMINATION OF THE HUDSON RIVER ECOSYSTEM

COMPILATION OF CONTAMINATION DATA THROUGH 2008

HUDSON RIVER NATURAL RESOURCE DAMAGE ASSESSMENT

HUDSON RIVER NATURAL RESOURCE TRUSTEES

STATE OF NEW YORK

U.S. DEPARTMENT OF COMMERCE

U.S. DEPARTMENT OF THE INTERIOR

JANUARY 2013

Available from:
U.S. Department of Commerce
National Oceanic and Atmospheric Administration
Hudson River NRDA, Lead Administrative Trustee
Damage Assessment Center, N/ORR31
1305 East-West Highway, Rm 10219
Silver Spring, MD 20910-3281



HUDSON RIVER

**PCB CONTAMINATION OF THE HUDSON RIVER ECOSYSTEM
COMPILATION OF CONTAMINATION DATA THROUGH 2008**

EXECUTIVE SUMMARY

The Hudson River Natural Resource Trustees are conducting a natural resource damage assessment (NRDA) to investigate natural resource injuries that may have occurred due to the release of polychlorinated biphenyls (PCBs) from General Electric (GE) facilities at Hudson Falls and Fort Edward, NY. This report summarizes available information on PCB contamination in the Hudson River ecosystem, including historic information, but focusing particularly on data collected and analyzed between 2002 and 2008 as part of ongoing NRDA activities. The Hudson River, for greater than 200 miles below Hudson Falls, NY, is extensively contaminated with PCBs. Surface waters, sediments, floodplain soils, fish, birds, wildlife, and other biota are all contaminated with PCBs. PCB concentrations are generally highest in those areas that are closer to the GE facilities, which are responsible for the majority of the area's PCB contamination. PCB concentrations tend to decrease with increasing distance downstream from these facilities. PCB concentrations upstream of the plant sites are substantially lower than the levels downstream.

This report also compares PCB concentrations in environmental media to regulatory standards and guidance criteria as well as to effects thresholds from the scientific literature. The more frequently these levels are exceeded, and the greater the magnitude of the exceedance, the more likely it is that PCBs have injured the natural resource(s) in question. Examples of exceedances described in this report include, but are not limited to:

- In water, exceedances of water quality standards and criteria;
- In sediments, exceedances of adverse effects levels for benthic organisms;
- In fish, exceedances of the U.S. Food and Drug Administration's (FDA) tolerance level for edible portions of fish;
- In mink, exceedances of levels associated with reproductive impairment;
- In snapping turtles, exceedances of levels associated with the latent mortality in juveniles;
- In bullfrogs, exceedances of levels associated with ecologically significant adverse effects (metamorph malformations and altered sex ratios); and
- In birds, exceedances of levels associated with reproductive impairment.

The frequency and severity of these exceedances varies by location and date; however, there are numerous instances in which the measured PCB concentrations exceed the selected benchmark by a factor of 10, 100, or more.

The information in this report demonstrates that high levels of PCB contamination have existed for decades in the Hudson River ecosystem. The data also suggest that PCBs are likely to be causing serious adverse effects to the area's biota.

The Hudson River Natural Resource Trustees' (HRNRT) natural resource damage assessment (NRDA) for the Hudson River ecosystem is ongoing. Additional information about the NRDA can be found on the following websites:

<http://www.darp.noaa.gov/northeast/hudson/index.html>

<http://www.dec.ny.gov/lands/25609.html>

<http://www.fws.gov/contaminants/restorationplans/HudsonRiver/index.html>

TABLE OF CONTENTS

EXECUTIVE SUMMARY

CHAPTER 1: INTRODUCTION..... 1

CHAPTER 2: PCB CONTAMINATION IN THE HUDSON RIVER..... 5

RIVER WATER..... 5

SEDIMENTS..... 7

GROUNDWATER..... 7

ADULT AQUATIC INSECTS..... 9

FISH 9

FLOODPLAIN SOILS 14

SMALL TERRESTRIAL MAMMALS AND THEIR PREY..... 14

TURTLES AND FROGS 16

BIRDS 18

BATS 20

MINK AND OTTER..... 20

CHAPTER 3: CONCLUSIONS..... 23

FURTHER INFORMATION..... 24

REFERENCES 25

LIST OF EXHIBITS

Exhibit 1	Bioaccumulation of PCBs.....	2
Exhibit 2	The General Electric Hudson Falls Plant on the Hudson River.....	3
Exhibit 3	Hudson River Below Corinth, New York.....	4
Exhibit 4	PCBs in Hudson River Surface Waters, 1975-2007.....	5
Exhibit 5	PCBs in Hudson River Surface Water, All Locations, 1977-2007.....	6
Exhibit 6	PCB Concentrations in Hudson River Mainstem Surface (≤ 12 inches) Sediments by River Section, 1975-2007.....	8
Exhibit 7	PCBs in Insects Emerging from the Hudson River, 1998.....	9
Exhibit 8	Total PCBs (ppm) in Fillets of Fish Caught in the Thompson Island Pool (River Section 1) between 2004 and 2008.....	10
Exhibit 9	Hudson River Fish Consumption Advisories for 2008-2009.....	12
Exhibit 10	Total PCBs (ppm) in Whole Fish Caught South of the General Electric Plants, 1977-2008.....	13
Exhibit 11	Upper Hudson River Floodplain.....	14
Exhibit 12	Percent of Samples by PCB Range (ppm) in Surficial (≤ 6 Inches Deep) Soils Associated with the Upper Hudson River, 1998-2008.....	15
Exhibit 13	PCBs (ppm) in Earthworms and Small Mammals from the Upper Hudson Floodplain, 2000-2001.....	15
Exhibit 14	PCBs in Snapping Turtle Eggs from the Hudson River Vicinity, 2002.....	16
Exhibit 15	PCBs in Bullfrog Tadpoles from the Upper Hudson River, 2003.....	17
Exhibit 16	PCBs in Bird Eggs Collected Near the Hudson River, 1994-2003.....	19
Exhibit 17	PCBs in Hudson River Waterfowl Fat, 1981-1984.....	20
Exhibit 18	PCBs (ppm) in Mink Livers as a Function of Stream Distance from the Upper Hudson River, 1989-2002.....	21

ACRONYMS AND ABBREVIATIONS

DOI	U.S. Department of the Interior
EPA	U.S. Environmental Protection Agency
FWS	U.S. Fish and Wildlife Service
fw	Fresh wet weight
GE	General Electric Company
HRNRT	Hudson River Natural Resource Trustees
NOAA	National Oceanic and Atmospheric Administration
NRDA	Natural Resource Damage Assessment
NYDEC	New York State Department of Environmental Conservation
PCBS	Polychlorinated biphenyls
ppb	Parts per billion (for example, one microgram per kilogram, or $\mu\text{g}/\text{kg}$)
ppm	Parts per million (for example, one milligram per kilogram, or mg/kg)
pptr	Parts per trillion (for example, one nanogram per kilogram, or ng/kg)
tPCBs	Total polychlorinated biphenyls
USGS	U.S. Geological Survey
ww	Wet weight

CHAPTER 1: INTRODUCTION

The Hudson River ecosystem below Hudson Falls, NY is extensively contaminated with PCBs. Federal and State Trustees are investigating injuries to living organisms and other natural resources that may have been caused by PCBs present in the Hudson River and surrounding environment.

The Trustee agencies are the U.S. Department of Commerce (DOC), the U.S. Department of the Interior (DOI), and the State of New York. These entities have each designated representatives that possess the technical knowledge and authority to perform natural resource damage assessments (NRDAs). For the Hudson River, the designees are the National Oceanic and Atmospheric Administration (NOAA), which represents the U.S. Department of Commerce, the U.S. Fish and Wildlife Service (FWS), which represents the concerned DOI bureaus (FWS and the National Park Service), and the New York State Department of Environmental Conservation (NYSDEC), which represents the State of New York.

This report provides an overview of PCB contamination of the Hudson River ecosystem including some historical information, but focusing particularly on data collected and analyzed between 2002 and 2008 as part of the ongoing natural resource damage assessment (NRDA). (Post-2008 data may be presented in an update to this report.) The Hudson River Natural Resource Trustees (HRNRT) are using this information together with the results of ongoing and future studies to assess injury to natural resources, and to determine the amount and type of restoration necessary to compensate the public for natural resource injuries and associated lost services.


PCBs, or polychlorinated biphenyls, are a group of highly toxic compounds that are known to cause cancer, birth defects, reproductive dysfunction, growth impairment, behavioral changes, hormonal imbalances, damage to the developing brain, and increased susceptibility to disease in animals (HRTC 2002).

PCBs accumulate in living organisms (Exhibit 1). They are transmitted from animal to animal via the food chain, and from parent to offspring in eggs. PCBs are stored in body fat, and the PCBs that have accumulated in a mother's body can be passed to early life stages via eggs (e.g., Kelly et al. 2008), and through placental and lactational exposures (e.g., Bleavins et al. 1981). PCBs can also contaminate the air, entering organisms via the lungs or gills, and are absorbed through the skin, such as from contact with contaminated soil (ATSDR 2000).

PCB concentration measurements in sediments, soils, water, and biota are frequently reported in units of parts per million (ppm), parts per billion (ppb), or even parts per trillion (ppt), depending on the material measured and on the amount of PCBs present in the material. A PCB concentration of one part per million (or billion, or trillion) for a sample means that the sample contains one part of PCBs per million (or billion, or trillion) parts of whatever is being analyzed (sediments, soil, water, animal tissue or other material).

Two General Electric facilities (Exhibit 2) have been identified as the predominant historical sources of PCBs to the Hudson River. EPA has estimated that these plants released up to 1,300,000 pounds of PCBs to the Hudson River between the 1940s and 1977 (EPA 2002). After 1977, seepage of PCBs from the bedrock beneath these General Electric facilities, combined with erosion of remnant deposits and contaminated banks, continued to release PCBs to the river for some time (EPA 2000a). In more recent years, the amount of PCBs released from seepage has been markedly reduced due to the continuing performance of remedial measures at these facilities (NYSDEC 2004).

EXHIBIT 1: BIOACCUMULATION OF PCBs



Source: Adapted from Wisconsin Department of Natural Resources. Used with permission.
<http://dnr.wi.gov/org/water/wm/foxriver/whatarepcbs.html>.

EXHIBIT 2: THE GENERAL ELECTRIC HUDSON FALLS PLANT ON THE HUDSON RIVER

Source: <http://www.fws.gov/contaminants/images/gehudsonriverepaimage.jpg> (EPA photo)

The General Electric Hudson Falls plant used PCBs for manufacturing operations beginning in 1947. The abandoned Allen Mill structure situated just below the Hudson Falls Plant failed in 1991, leading to a temporary spike in outflow of PCBs to the Hudson River.

In general, PCB concentrations in the Hudson River ecosystem have been highest immediately downstream of the General Electric (GE) facilities. The GE facilities are located in Hudson Falls and Fort Edward (Exhibit 3), upstream of the Thompson Island Pool also known as River Section 1. River Section 1 is about six miles long, extending from the location of the former Fort Edward Dam downstream to the Thompson Island Dam.

The adjacent downstream portion of the river is River Section 2, consisting of the Fort Miller Pool immediately below the Thompson Island Dam and the Northumberland Pool, which is connected to the Thompson Island Pool by the

land cut. River Section 2 is about five miles long and extends from the Thompson Island Dam to Northumberland Dam. River Section 3 is much larger – about 29 miles long—and extends from the Northumberland Dam downstream to the Federal Dam at Troy and is comprised of several pools separated by a series of locks. Collectively, these three river sections, approximately 40 miles in combined length, comprise the Upper Hudson. The Lower Hudson consists of the portion of the river south of the Federal Dam downstream to the Battery in New York City. This section of the river is tidally influenced and is over 150 miles long.

The Hudson River at and below Hudson Falls, New York, has been contaminated extensively by PCBs from GE facilities. The PCBs present a serious and long-term threat to the health of the Hudson River ecosystem and pose a potential health threat to people who eat fish or who eat other organisms that inhabit the river and the surrounding area. Numerous studies have documented PCB contamination in the surface water, groundwater, sediments and floodplain soils of the Hudson River, as well as in living resources at every level of the Hudson's aquatic, terrestrial, and wetland-based food chains.



CHAPTER 2: PCB CONTAMINATION IN THE HUDSON RIVER

RIVER WATER

Water sampling has occurred throughout the river south of Hudson Falls; however the large majority of samples were taken from the Upper Hudson. Since PCB measurements of Hudson River water began in the 1970s, PCB levels in the river below Hudson Falls have routinely exceeded state and Federal water quality criteria developed to protect living organisms. Over 80 percent of over 6,000 water samples tested since 1975, from Hudson Falls to the Battery in Manhattan have contained PCBs at levels 10 to 10,000 times higher than that deemed safe for aquatic life, fish-eating wildlife and human consumers of fish (HRNRT 2008d, see Exhibits 4 and 5). Furthermore, this percentage very likely underestimates the extent of past contamination in the Hudson because early tests for PCBs were not as sensitive as more modern methods.

PCB levels in the river have also exceeded the current drinking water standard (0.09 ppb) in about two percent of samples taken from applicable locations (HRNRT 2008d). This standard was promulgated on March 12, 1998. The previous drinking water standard, promulgated in 1985, was 0.01 ppb, and nearly 80 percent of samples in applicable locations exceeded that value. Of note, drinking water standards apply only to portions of the river with certain designated water classes, which are located approximately between river miles 162 and 156 (the confluence with the Mohawk River to the Lock 2 Dam), and between river miles 129 to 65 (Chelsea to Houghtaling Island) (HRNRT 2008d).

TOXICITY OF PCBs

PCBs are highly toxic—even very small amounts are considered hazardous. The New York State water quality standard for protection of human consumers of fish is 0.001 parts per trillion, or just 1/1,000th of a drop of PCBs in one trillion (1,000,000,000,000) drops of water.

The U.S. Environmental Protection Agency (EPA) issued the first regulatory standard for PCB levels in surface water in 1977, determining that PCBs in water at levels as low as 1 part per trillion (ppt) pose an unacceptable risk to humans and aquatic organisms. As scientific understanding of PCBs grew and the ability to measure PCB concentrations improved, EPA and the State of New York established criteria for PCBs in surface water to protect specific groups of organisms. The current criteria set levels for protection of salt water aquatic life (30 ppt on a total PCB basis) and freshwater aquatic life (14 ppt on a total PCB basis). Current regulatory standards are also in place for fish-eating wildlife (0.12 ppt on a total PCB basis) and human consumers of fish (0.001 ppt on a total PCB basis). PCB levels in the Hudson have far exceeded these amounts in the vast majority of water samples tested since the 1970s, with levels detected ranging to over 38,000 parts per trillion in the Upper Hudson. In 2008, the Trustees determined that the Hudson River's surface water has been, and continues to be, injured as a consequence of PCB exposure (HRNRT 2008d).

EXHIBIT 4: PCBs IN HUDSON RIVER SURFACE WATERS, 1975-2007





EXHIBIT 5: PCBs IN HUDSON RIVER SURFACE WATERS, ALL LOCATIONS, 1975-2007





Data sources: Measurements taken by the U.S. Geological Survey, which in 1975 initiated regular monitoring of PCBs in the water column at Waterford and then expanded its monitoring program to a total of seven stations, all within the Upper Hudson. Additional data sources include measurements taken by GE, whose sampling program began in 1989, and which encompasses 124 locations, 120 of which are located in the Upper River, and smaller datasets provided by EPA (2000b), Litten (2003), and Bopp et al. (1985), all three of which included some sampling in both the Lower and Upper Hudson.

Notes: Data are shown for all years (1977-2007) and for the period after 3/12/1998 to 2007. The depicted 1 ppt EPA guidance criterion was established in 1976, while the 0.12 ppt and 0.001 ppt New York State water quality standards were promulgated on 3/12/1998. Because water quality criteria apply only after their issuance, only measurements taken after availability of the relevant criteria are included in this figure. The presented data also reflect only those samples in which PCBs were detected (about 80% of all measurements).

ABOUT THE GRAPHS IN THIS REPORT

Graphs in this report generally are presented in two styles. Where data are relatively few, individual measurements are shown. Where data are more numerous, a box-and-whisker style is used to illustrate the general distribution of the values. The whiskers represent the minimum and maximum values. The top and bottom of the colored box represent the 75th and 25th percentiles, respectively, and the central line represents the median value.

In addition, because of the wide range in PCB concentrations depicted in many of the graphs in this report, the information is shown in logarithmic scale to allow all the data to appear visible on the same graph. On a logarithmic scale, every ten-fold difference (such as between 1 and 10, or between 10 and 100) is depicted as an equally-sized interval. The sample figures below illustrate the visual difference between the same sample dataset plotted on both a linear and a logarithmic scale.



It is not possible to show values of zero on a logarithmic scale; therefore, samples in which PCBs are not detected (and which are assumed for purposes of this report to contain no PCBs) are discussed in the figure legends.

SEDIMENTS

Over the past 60 years, large quantities of PCBs from the Hudson River have settled out into the riverbed over a distance spanning more than 200 miles downstream of Hudson Falls to New York Harbor and beyond. Surface sediment (considered for purposes of this report to be sediments 12 or fewer inches deep) PCB concentrations are significantly higher in the Upper Hudson than the Lower Hudson. While surface concentrations in River Sections 1 and 2 are similar, concentrations decline in River Section 3, and decline further in the estuary. Through 2008 (the time period that is the focus of this report), PCB measurements in surface sediments of the Upper Hudson and some parts of the Lower Hudson have been well in excess of levels associated with toxic impacts (Exhibit 6).

PCBs in the riverbed continue to contaminate aquatic insects, mussels and other invertebrates that live in the sediment. During sampling conducted by GE in the Upper Hudson between 2002 and 2007, PCBs were detected at levels as high as 1,650 ppm in surficial sediments (top 12 inches), and this particular value was measured within the top two inches. This level of contamination creates a hazardous environment for exposed biota: for instance, NYSDEC (1999) developed sediment-based PCB screening criteria of 0.042 ppm for wildlife bioaccumulation, 0.58 ppm for chronic benthic effects, and 83 ppm for acute benthic effects.¹ Sediments with concentrations above these levels “are considered to be contaminated, and [are] potentially causing harmful impacts to marine and aquatic ecosystems” (ibid.). Approximately 99% of surficial (\leq 12 inches deep) remedial design samples exceed the NYSDEC 0.042 ppm criterion, while about 97% and 17% of these samples exceed the 0.58 ppm and the 83 ppm criteria, respectively.²

In addition, EPA determined that sediment concentrations in excess of 3 ppm pose a risk to amphibians in the Housatonic River,³ another site that has been contaminated with PCBs from GE facilities (Weston Solutions 2003). This level is exceeded in over 70% of the Hudson River surficial sediment samples collected as part of the remedial design sampling. Follow-up studies on sediment are currently underway (HRNRT 2008b).

GROUNDWATER

Groundwater in the vicinity of GE’s Hudson Falls and Fort Edward plants is heavily contaminated with PCBs, along with high concentrations of volatile and semi-volatile organic compounds. When the extent of contaminated groundwater in this area was discovered in the 1980s, the Town of Fort Edward issued bonds to pay for construction of a water system to serve Fort Edward Water District No. 1.




In the early 1990s, New York State determined that the releases of PCBs from the GE Hudson Falls and Fort Edward plant sites, including the migration of contaminated groundwater from beneath the Hudson Falls plant, represented a significant ongoing source of PCBs to the Hudson River. Implementation of remedial actions at both the Hudson Falls and Fort Edward capacitor plant sites since the early 1990s has resulted in marked reductions of PCBs released to the river from the plant sites. Remedial work is continuing at both plant sites.⁴

¹ NYSDEC (1999) presents values on a $\mu\text{g/g}$ organic carbon (OC) basis (i.e., 1.4 $\mu\text{g/g}$ OC, 19.3 $\mu\text{g/g}$ OC, and 2760.8 $\mu\text{g/g}$ OC). The presented criteria have been converted into a mg/kg dry weight basis assuming that sediments contain 3% organic carbon, the average value for the Hudson River.

² The presented percent exceedance figures are calculated comparing the original threshold values (in $\mu\text{g/g}$ OC) to sediment concentration values expressed in the same units. Samples lacking an organic carbon measurement are not included in these calculations.

³ This value was selected as the “maximum acceptable threshold concentration” although adverse effects (including high mortality and developmental delays) were exhibited by leopard frog larvae exposed to all Housatonic site sediments. These sediments had PCB concentrations as low as 0.15 ppm (Weston Solutions 2003).


EXHIBIT 6: PCB CONCENTRATIONS IN HUDSON RIVER MAINSTEM SURFACE (≤ 12 INCHES) SEDIMENTS BY RIVER SECTION, 1975-2007



Data source: NOAA (2010).

Sediment data are grouped by data-generation project. This roughly approximates time periods and also reflects the different sampling strategies employed. Non-detects (less than 1% of the samples) are not included in this analysis and lab/field duplicates have also been removed, as have samples not taken in the river's mainstem. Surface sediments samples are defined as those that begin at the surface and have a lower depth of 12 inches or less. NYSDEC (1999) developed the depicted criteria for wildlife accumulation (1.4 µg/g OC), chronic benthic toxicity (19.3 µg/g OC), and acute benthic toxicity (2760.8 µg/g OC). Assuming 3% organic carbon in sediments (average for the Hudson River), these values become 0.042, 0.58, and 83 mg/kg sediment.

EXHIBIT 7: PCBs IN INSECTS EMERGING FROM THE HUDSON RIVER, 1998



Data source: HRNRT (2009a), NOAA (2010).

Notes: These insects inhabit the Hudson riverbed during their larval stage. As adults, they can fly and thereby provide a potential pathway for contamination of the floodplain food web. All samples contained detectable concentrations of PCBs. Each value represents a composite sample.

The New York Department of Health determined that the Village of Stillwater Well Field, in Saratoga County, is a groundwater resource that is under the influence of the surface water of the Hudson River. Results of sampling of several of the wells in the Stillwater Well Field shows that these groundwater resources exceed New York State’s PCB standard of 0.09 ppm in fresh groundwater (Malcolm Pirnie 2009). As a result, the Village of Stillwater recently has paid for construction of a pipeline in order to change its water source to the Saratoga County Water Authority (Cignoll 2012).

ADULT AQUATIC INSECTS


Hudson River fish, birds, and wildlife can also become contaminated with PCBs through consumption of aquatic insects such as dragonflies, mosquitoes and mayflies, which live in or on the river’s bottom as larvae but emerge from the river as flying insects in their adult forms. A 1998 study found PCB levels in these emerging adult insects along the banks of the Upper Hudson to be as high as 6.3 ppm (HRNRT 2009a). The study also found that PCB levels in insects captured in the Upper Hudson were over ten-fold higher than levels in insects captured in the Lower Hudson, about 90 miles downstream of the Troy Dam (Exhibit 7).

FISH

Hudson River fish downstream of the GE plants have been contaminated by PCBs. Fish not only absorb PCBs directly from the river water but are also exposed through the ingestion of contaminated prey, such as insects, crayfish, and smaller fish. In addition, fish, especially eggs, can be exposed to contaminants through river sediments. Contaminated fish can, in turn, be eaten by birds, wildlife or even humans, and can serve to expose these groups to PCBs as well.

In the 1970s, several Hudson River largemouth bass fillets were contaminated at levels in excess of 2,000 ppm (NOAA 2010). After 1977, when GE stopped direct discharges of PCBs to the river, PCB levels in Hudson River fish dropped considerably but since the early 1980s have generally remained stable⁵ at relatively high levels. Sampling results indicate that PCB concentrations in fish tend to be highest in the Upper Hudson downstream of the GE plants at Hudson Falls and Fort Edward, and generally decline with increasing distance down the river, with a less pronounced gradient in the Lower Hudson.

PCBs in Hudson River fish may pose significant health risks to human consumers of fish. The U.S. Food and Drug Administration does not allow the commercial sale of fish for consumption by humans where PCB levels exceed 2 ppm, and the Environmental Protection Agency’s remedial action objective (RAO) for the protection of human health is 0.05 ppm in fish fillets, based on an adult consumption rate of one half pound meal per week (EPA 2000a). In fish collected since 2000 in the Upper Hudson, this level was exceeded in approximately 75-90% of largemouth bass, smallmouth bass, brown bullhead, and yellow perch fillet samples, depending on the species. In fact, about 40-70% of these fillets (depending on species) exceeded this level by a factor of ten or more. (See Exhibit 8 for a subset of these data.) In striped bass and white perch fillets collected in the Lower Hudson over the same time period, EPA’s 0.05 ppm RAO was exceeded in over 95% of samples, and about 65% of these samples exceeded this level by a factor of ten or more.



⁵ One exception to the relative stability of PCB levels in fish in the Hudson was the spike in levels that occurred in 1992 and 1993 following a release of PCBs resulting from the collapse of a structure at the Allen Mill, an abandoned industrial building adjacent to GE’s plant at Hudson Falls. Concentrations dropped through the 1980s until the Allen Mill event, increased, then decreased after the spike. The rate of decline was generally greater prior to, rather than after the Allen Mill event. After recovery from the event, the rate of decline remained relatively stable, especially for sampling locations nearest the plant sites (Sloan et al. 2005).

Because of excessive levels of PCBs in Hudson River fish, New York State sharply limited fishing in the Hudson in 1976, closing most of the river's commercial fisheries, issuing an "eat none" advisory and prohibiting all recreational angling throughout the 40 miles of the Upper Hudson downstream of Bakers Falls. Most of the commercial closures remain in place to this day. The "eat none" advisory and the restriction on taking fish for this section of the Upper Hudson has been in place for 36 years. The ban on recreational angling in the Upper Hudson remained in place until 1995, when the State modified the regulations to permit catch and release fishing within that reach. Consumption advisories have also been established throughout the Lower Hudson River; these have varied over time by species and location.


The 2008-2009 advisories (Exhibit 9)⁶ include a "no consumption" advisory for all fish taken from the river between South Glens Falls (upstream of Hudson Falls) and the Troy Dam. "No consumption" advisories also exist for the general population for most species of fish from the Hudson River below the Troy Dam as far as Catskill, and advisories are in place restricting consumption to one meal per month for many species between Catskill and the Battery. In addition, women of childbearing age and children under age 15 are advised not to eat fish or crabs taken from the length of the river from Hudson Falls to the Battery.

Beyond the risks to human consumers of fish, studies at other contaminated sites and extensive laboratory testing have shown that PCBs can cause a wide range of toxic effects to the fish themselves. These include increased susceptibility to disease, feminization of males, growth of cancerous tumors, reduced egg survival rates, and impairment and death of newly hatched fish. Skeletal deformities and organ hemorrhaging have also been found in fish exposed to PCBs, as well as hormonal disturbances and biochemical changes (HRTC 2002).

Sampling in the Upper Hudson River between 2000 and 2008 found PCB levels in whole fish up to 470 ppm, with the maximum value from earlier years reaching 571 ppm (Exhibit 10). For comparison, the scientific literature documents biochemical changes in fish with PCB levels less than 1 ppm (Niimi 1996). Also, research has found that PCBs can severely affect reproduction in the barbel, a species of fish in the same biological family as Hudson River carp and minnows. Barbels with whole body PCB levels of approximately 0.8 ppm wet weight (ww) experienced an excess egg mortality of 20% (Hugla and Thomé 1999). Follow-up studies on fish are currently underway (HRNRT 2001, 2009b).


⁶ These advisories are referenced because of the focus of this report on data through 2008. Current advisories are available in NYSDOH (2011).

EXHIBIT 9: HUDSON RIVER FISH CONSUMPTION ADVISORIES FOR 2008-2009



Data Source: NYSDOH (2008).

EXHIBIT 10: TOTAL PCBs (PPM) IN WHOLE FISH CAUGHT SOUTH OF THE GENERAL ELECTRIC PLANTS, 1977-2008



Data Source: NOAA (2010)

Notes: Data are presented for the 10 species of Hudson River fish with the highest number of whole body samples. Samples identified as duplicates and non-detects (less than 2%) are not included in the analysis. Some records represent composites of multiple fish such that the listed number of values is less than the number of fish caught and analyzed. Fish data are broken into the following time periods: 1977-1980; 1981-1990 (passage of CERCLA through just before the Allen Mill event); 1991-1996 (representing the immediate post-Allen Mill period); 1997-2008 (up through the time just before Phase 1 dredging began). Data are compared to the level at which egg mortality has been observed to increase by 20% over control levels in barbels (a fish in the same family as Hudson River carp and minnows, Hugla and Thomé 1999). The 0.8 ppm wet weight value reflects the paper's 0.40 µg/g dry weight egg threshold converted into whole body wet weight using an egg-to-adult PCB dry weight ratio (calculated from Table 2 in the Hugla and Thomé 1999), assuming 80 percent moisture. Note: additional data for fish filets is available, but is not depicted in the exhibit above.

FLOODPLAIN SOILS

The Hudson River floodplain is periodically inundated with water and sediments from the river (Exhibit 11). This inundation can result in the deposition of PCBs onto the river's banks as well as onto the adjoining wetlands and forested floodplain. This contaminated material is ingested by small animals such as insects and worms, which are a source of food for other animals. Because PCBs accumulate in the fatty tissues of animals, each time an animal is exposed to PCBs, the PCB level in its body can rise, leading to increased potential for toxic effects. As animals higher on the food chain consume other animals, PCBs can become more concentrated, potentially leading to very high levels of these toxic chemicals in larger animals such as mammals, birds, and humans.

Several studies have measured PCB concentrations in the Upper Hudson floodplain soils. Concentrations in surficial soils (considered for purposes of this report to be soils six or fewer inches deep) range up to 1,040 ppm in River Section 1, up to 358 ppm in River Section 2, and up to 63 ppm in River Section 3 (NOAA 2010). Some studies found that PCB concentrations in floodplain soils tend to be highest in soils closer to the river (SEA 2002, Weston Solutions 2005), and closer to the GE plants, with concentrations decreasing downstream (SEA 2002). Exhibit 12 presents distributional information on PCB concentrations in surficial (≤ 6 inches deep) soils, grouped first by study source and secondarily by river section. Of note, the data presented represent a range of distances from the river and have their origin in collection efforts that utilized different sampling strategies.

For comparison, Efroymsen et al. (1997) developed a preliminary remediation goal (PRG) for total PCBs (tPCBs) in soils of 0.371 ppm. This level generally "correspond[s] to small effects on individual organisms which would be expected to cause minimal effects on populations and communities" of wildlife. The PRGs may not be sufficiently protective of species of special concern which are based on effects on individual organisms and should be based on no-observed adverse-effects levels." Of the datasets shown in Exhibit 12, over a third of samples in River Sections 1 and 2 exceed this value, and approximately a fifth of samples in River Section 3 exceed this value.

SMALL TERRESTRIAL MAMMALS AND THEIR PREY

PCBs from contaminated floodplain soils along the Hudson River have entered the terrestrial food chain. Earthworms collected in 2000 from the Upper Hudson River floodplain contained PCBs at levels averaging 7.7 ppm and as high as 23.9 ppm (Exhibit 13).


Shrews, mice, and meadow voles are important prey items for larger animals such as mink and raptors. Their contamination levels suggest that these small mammals may be an important pathway for PCB exposure to their predators.

EXHIBIT 11 : HUDSON RIVER FLOODPLAIN



Source: Joseph Steinbacher, Versar, Inc.


EXHIBIT 12: PERCENT OF SAMPLES BY PCB RANGE (PPM) IN SURFICIAL (≤ 6 INCHES DEEP) SOILS ASSOCIATED WITH THE UPPER HUDSON RIVER, 1998-2008



Data source: NOAA (2010)

Notes: ND = non-detect. Samples identified as duplicates and apparent field replicates have been eliminated from the analysis. Samples have a lower depth of not more than 6 inches. Results are grouped by study effort(s). Of note, Efraymson et al. (1997) developed a PRG for total PCBs of 0.371ppm, a level that generally “correspond[s] to small effects on individual organisms which would be expected to cause minimal effects on populations and communities.”

EXHIBIT 13: PCBs (PPM) IN EARTHWORMS AND SMALL MAMMALS FROM THE UPPER HUDSON FLOODPLAIN, 2000-2001



Data sources: SEA (2002), HRNRT (2010), NOAA (2010). All samples contained detectable concentrations of PCBs. The mouse and vole values each represent composites of four to five animals, whereas the shrew data represent individual animals.



TURTLES AND FROGS

Reptiles from the Hudson River are also contaminated with PCBs. Snapping turtles collected in 1998 and 2000 contained very high levels of PCBs, ranging to over 3,000 parts per million in fat (NOAA 2010).

PCBs have been associated with behavioral abnormalities and changes to biochemistry in adult snapping turtles (HRNRT 2002), and PCBs are also passed from adult female turtles to their eggs. Snapping turtle eggs collected in 2002 contained elevated PCB levels, ranging up to 31.8 ppm (HRNRT 2005a, Exhibit 14). PCBs in snapping turtle eggs have also been linked to latent mortality: Eisenreich et al. (2009) found that snapping turtles hatched from Upper Hudson River PCBs eggs suffered a 60 percent mortality rate through 14 months of age, compared with a 10 percent rate for animals hatched from reference area eggs. Furthermore, the mortality rate was correlated with total PCBs in the collected eggs: the authors calculated a relationship between PCB egg concentrations and mortality, which suggests that levels of approximately 3.3 ppm decrease survival to about 80% of what it would have been absent PCB exposure (Eisenreich et al. 2009). Turtle eggs are also a pathway for PCB contamination to animals that consume these eggs, such as other reptiles, birds, and mammals, and potentially humans.

New York State has a long-standing statewide health advisory recommending no consumption of snapping turtles (or soups made with their meat) by women of childbearing age and children under the age of 15. The advisory further recommends that all others carefully trim all fat, and discard fat, liver, and eggs prior to cooking, to reduce exposure to contaminants. In 1993-1994, the advisory was clarified to explain that PCBs represent the chemical of concern (NYSDOH 1993). Concern about PCBs in turtles predates this clarification, however: since 1979, the elevated levels of PCBs in turtles—including Hudson River turtles—have been a driving concern in the state’s warnings to citizens to avoid consumption of these animals (Funicello 1979).

EXHIBIT 14: PCBs IN SNAPPING TURTLE EGGS FROM THE HUDSON RIVER VICINITY, 2002




Data sources: HRNRT (2005a) and NOAA (2010).

Notes: PCBs were detected in all samples. Each sample represents a composite of three to five eggs. “Regions” are defined as shown in this figure (from HRNRT (2005a)); these regions differ from the “river sections” discussed elsewhere in this report. The vertical lines within the sampling data represent median values. The 3.3 ppm value associated with 20% mortality is derived from Eisenreich et al. (2009).

Amphibians from the Hudson River are also contaminated with PCBs. Bullfrog tadpoles collected in 2003 had levels as high as 9.3 parts per million (HRNRT 2007b; also see Exhibit 15). For comparison, in the Housatonic River (another PCB-contaminated site), EPA determined that PCB concentrations of 1 ppm in wood frogs is the level at which “significant adverse effects begin to occur, and response became frequent and more severe at approximately 10 mg/kg” (Weston Solutions 2003). These responses include the frequency of malformations in metamorphs and male:female sex ratios of 0.7 or less.

EXHIBIT 15: PCBs IN BULLFROG TADPOLES FROM THE UPPER HUDSON RIVER, 2003



Data source: HRNRT (2007b), NOAA (2010).

Notes: PCBs were detected in all samples. Each sample represents a composite of one to 12 tadpoles (most commonly, two to six tadpoles) at a similar developmental stage. The “risk level” is the value determined by EPA to pose a significant risk to amphibians (Weston Solutions 2003). The vertical lines within the sampling data represent median values. The reference site locations are varied and are not depicted on the map.

In the Hudson, the levels and specific types of PCBs detected in bullfrog tadpoles closely mirror those found in the sediments where the tadpoles were living. This suggests that in addition to receiving a burden of PCBs from their parents in the egg, these tadpoles are acquiring PCBs from the river environment they inhabit. Furthermore, PCB levels in sediments from known amphibian breeding areas of the Hudson River are at ecologically significant levels, suggesting the potential for injury to these organisms (HRNRT 2008a). The Trustees are investigating additional options to assess amphibian injury (HRNRT 2008c).

BIRDS

Birds in the vicinity of the Hudson River have been exposed to PCBs from the fish, insects, and other animals in their diet, as well as through the soil they ingest while feeding. In studies at other locations and in the laboratory, PCBs have been linked to a wide range of adverse impacts to birds, including disease, behavioral abnormalities, genetic mutations, physical deformities, changes in brain chemistry, reduced hatching rates, mortality of embryos, and death of adult and juvenile birds (HRTC 2002). In addition, PCB-contaminated birds and bird eggs are a source of PCB contamination for the animals that consume them, such as reptiles, mammals, and other birds.

More than 150 species of birds inhabit the Hudson River region at various times of the year, including waterfowl, wading birds, shorebirds, songbirds, and raptors. Birds are an integral part of the ecosystem, playing a number of important ecological roles, including seed distribution, plant pollination, and insect control. Some birds are also important prey items for other animals.

Several studies have confirmed that birds and their eggs in the Hudson River region are contaminated with PCBs (Exhibit 16). Tree swallows, which eat insects that inhabit the river bottom as larvae, are particularly likely to accumulate PCBs. In the mid-1990s, tree swallows were found to contain extremely high levels of PCBs, ranging up to 62 parts per million in nestlings (McCarty and Secord 1999a) and 190 parts per million in adult swallows (HRNRT 2011b). These are among the highest PCB levels ever reported in tree swallows from any location (McCarty and Secord 1999a). Tree swallow eggs were also found to be highly contaminated with PCBs, containing levels up to 77 ppm (McCarty and Secord 1999a).⁷ In addition, studies conducted on tree swallows along the Upper Hudson during the 1990s found indications of disrupted reproductive functioning, including high incidence of nest abandonment (McCarty and Secord 1999a), inability to build normal nests (McCarty and Secord 1999b), and abnormal appearance of younger females during the breeding season (McCarty and Secord 2000). Similar signs have been associated with the effects of PCBs on hormone levels in other species (McCarty and Secord 1999b).

Elevated PCB levels were found in other birds in the vicinity of the Hudson during the mid- to late 1990s, including up to 78 ppm in eastern bluebird nestlings (HRNRT 2011b), 220 ppm in the fat of great blue heron nestlings (HRNRT 2011d), and 85.8 ppm in the fat of a bald eagle (HRNRT 2011b). Bald eagle eggs that failed to hatch were collected from the Lower Hudson in 1998-1999 and 2003 and 2004; these eggs were found to be highly contaminated with PCBs (HRNRT 2011c,d,e,f,g).


In 2002-2003, the eggs from several species of birds were collected from areas adjacent to the Hudson River (HNRTC 2005b,c). PCB levels as high as 56 ppm were detected, with the highest levels of contamination found in kingfisher and spotted sandpiper eggs (HRNRT 2005b). Harris and Elliott (2011) reviewed the effects of PCBs on wild birds and note that species exhibit varying sensitivity to PCBs. Among wild birds of “intermediate” (or possibly intermediate) sensitivity, critical egg thresholds for reproduction have ranged from 6 ppm to 50 ppm (Table 14.8, Harris and Elliott 2011). The lower level of 6 ppm has been exceeded in some eggs of many species of Hudson River birds, including the American robin, red-winged blackbird, common grackle, spotted sandpiper, Eastern phoebe, northern rough-winged swallow, tree swallow, Eastern screech owl, and belted kingfisher (Exhibit 16). Hudson River peregrine falcon eggs have also been shown to contain elevated levels of PCBs (HRNRT 2004a).

Additional reports have continued to confirm the presence of PCBs in Hudson River bird eggs of insectivorous, omnivorous and piscivorous bird species. Tree swallow eggs collected in 2004 contained an average of 6.8 ppm fww (Custer et al. 2010a); belted kingfisher eggs collected in 2004 contained an average 10.6 ppm fww (Custer et al. 2010c), and spotted sandpiper eggs collected in 2004 contained an average of 9.1 ppm fww (Custer et al. 2010b).⁸ Additional studies are currently underway to investigate the effect of PCBs on birds in the Hudson River region (HRNRT 2004b, 2005d, 2006b, 2007a and d, 2008e, 2009c).

⁷ Where available, the presented concentrations in eggs are fresh wet weight values (i.e., the values have been corrected to account for the duration of incubation). Where not available, values are in wet weight.

⁸ These publications report geometric rather than arithmetic means. Arithmetic means would likely be higher.

EXHIBIT 16: PCBs in Bird Eggs Collected Near the Hudson River, 1994-2003




Data sources: HRNRT (2005b,c), HRNRT (2011a), Secord and McCarty (1997).

Notes: All data shown are from 2002 and/or 2003 Trustee collections, except for tree swallow eggs, which were collected in 1994 and 1995. PCB values are in parts per million fresh wet weight. PCBs were detected in all samples. The presented critical threshold range is from Harris and Elliot (2011).

PCBs were also measured in waterfowl collected from hunters in New York State during the early 1980s (Kim et al. 1984, Kim et al. 1985). As shown in Exhibit 17, PCBs in the fat of a number of species of waterfowl have exceeded the 3 ppm U.S. Food and Drug Administration marketplace tolerance level for poultry (21 CFR 109.30). PCB concentrations ranged up to 22.7 parts per million in the fat of mallards, 124 ppm in a merganser, and 43 ppm in black ducks. These and other data showing similar patterns of contamination led the New York State Department of Health to issue a statewide waterfowl consumption advisory to protect human health. Further, limited data for mallards collected in 2000 confirm that PCB levels in these birds remain elevated, with up to 7.8 ppm found in fat (NOAA 2010). Further investigations of the level of PCBs in Hudson River waterfowl are underway (HRNRT 2008f).

EXHIBIT 17: PCBs in HUDSON RIVER WATERFOWL FAT, 1981-1984



BATS

Hudson River bats, which consume insects from both the floodplain and the aquatic ecosystem, also contain elevated levels of PCBs. A study conducted in 2001 and 2002 showed levels up to 0.64 ppm in the brains of big brown bats, and as high as 2.4 ppm in the brains of little brown bats (HRNRT 2007c).


MINK AND OTTER

Mink and otter of the Hudson River inhabit both the river and riverside habitats, coming into contact with (and ingesting) contaminated water, sediment, and soil as they build dens and forage for food. Mink and otter have also been exposed to PCBs through the fish, invertebrates, small mammals, and other prey they eat.

Mink are known to be sensitive to the effects of PCBs: jaw lesions in wild mink have been linked to PCB contamination at the Kalamazoo River Superfund Site (Beckett et al. 2005). These lesions have also been seen in captive mink kits fed diets containing 1 ppm PCBs (Bursian et al. 2006). PCBs can also have toxic effects on mink reproduction, causing reduced growth and increased mortality of offspring. Previous studies have found reduced kit growth and/or survival at dietary PCB concentrations of 1 ppm or less (Restum et al. 1998, Heaton et al. 1995), and a recent study using Hudson River fish found a dietary LC20 concentration⁹ for 6-week kit survivability, to be 0.34 ppm PCBs (Bursian et al. 2011). Given these data, the PCB contamination in the Hudson River environment appears to pose a high risk for mink: mink consume small mammals, fish, amphibians, reptiles, and birds, and as described previously, these organisms (see Exhibits 8, 10, 13,14, 15, 16) have frequently contained PCB levels in substantial excess of this level.

Measured concentrations of PCBs in Hudson River mink livers also suggest past and ongoing risks to mink (Exhibit 18). PCB levels in mink livers from the Hudson River watershed ranged up to 5.6 ppm, while Heaton et al. (1995) found PCB levels in livers of about 2 ppm to be sufficient to impair reproduction. Eight of the 33 mink (about 24%) caught within approximately one home range¹⁰ (6 km) of the Hudson River between 1998 and 2002 had liver PCB concentrations that exceeded this value.

EXHIBIT 18: PCBs (PPM) IN MINK LIVERS AS A FUNCTION OF STREAM DISTANCE FROM THE UPPER HUDSON RIVER, 1989-2002



Data source: D. Mayack, New York State Department of Environmental Conservation. A subset of these data is available in NOAA (2010). Notes: The vast majority of mink were caught adjacent to streams. Stream distance to the Hudson is calculated as the stream length (not the straight-line distance) between the collection location and the Hudson River. Except for three samples, all data depicted represent animals collected between 1998 and 2000. This figure uses an estimated home range for mink of 6 km (D. Mayack, personal communication). PCBs were detected in all samples. The liver threshold of 0.8 ppm represents a site-specific LC20 value for kit survivability from the Hudson River mink feeding study (Bursian et al. 2011).

⁹ The LC20 refers to the lethal concentration (LC) of a substance that is associated with a 20 percent mortality rate.

¹⁰ The term “home range” refers to the area that an animal normally uses throughout its life.

Data from the early 1980s, while few, suggest that mink exposure to PCBs was of a similar magnitude: total PCBs measured in seven mink livers from the upper Hudson River watershed ranged up to 1.7 ppm, with a median value of 0.5 ppm (Foley et al. 1988, R. Foley, personal communication). In the 1998-2002 collection of 33 mink caught within a home range of the Upper Hudson, liver concentrations of PCBs ranged up to 5.6 ppm, with a median value of 0.41 ppm.

A trapping study in the Upper Hudson floodplain during 1999 and 2000 found evidence of lower numbers of mink in areas closer to the Hudson, trapping an average of only 3.5 mink per 1,000 trap nights, compared to an average of 26.2 mink trapped in the same amount of time in upstream and distant sites (Mayack and Loukmas 2001). Together with extensive data from both laboratory tests and field studies at other contaminated sites linking PCBs to failed reproduction, these numbers suggest that elevated PCB levels in Hudson floodplain mink may be affecting survival and/or reproduction. Follow-up studies are currently underway (HRNRT 2006a, HRNRT 2011h, 2011i).

Otters may also be at risk from elevated PCB concentrations in the Hudson River watershed. Concentrations of PCBs in the livers of 31 otters caught within a home range (30 km) of the Hudson River between 1997 and 2002 ranged up to 22.5 ppm, with a median value of about 1.2 ppm.¹¹ Levels of about 0.63 ppm are believed to be deleterious to liver functioning in this species (Smit et al. 1996).¹²

¹¹ Data from D. Mayack, New York State Department of Environmental Conservation. A subset of these data are available in NOAA (2010).

¹² Above this concentration, Smit et al. (1996) estimate that 90 percent of animals would have reduced vitamin A levels in the liver. The 0.63 ppm value reflects the paper's estimated 21 µg/g lipid converted into the wet weight equivalent, assuming 3 percent lipids.

CHAPTER 3: CONCLUSIONS

The Hudson River, at and below Hudson Falls, New York, has been contaminated extensively by PCBs from GE facilities. Further, the resulting high levels of PCB contamination have existed for decades, and continue to exist, in the Hudson River ecosystem.

PCBs have contaminated the surface water, groundwater, sediments and floodplain soils of the Hudson River. Concentrations of PCBs in these environmental media exceed regulatory standards and criteria for their quality and use. Such exceedances include the following:

- Criteria for surface water quality are exceeded. The Hudson River's surface water has been, and continues to be, injured from PCB exposure. Additionally, groundwater around the GE facilities is heavily contaminated with PCBs and high concentrations of volatile and semi-volatile organic compounds.
- Edible portions of fish exceed the U.S. Food and Drug Administration's (FDA) tolerance level for PCBs, and there are advisories on fish consumption due to PCBs throughout the Upper and Lower Hudson Rivers.
- Consumption advisories are also in place for waterfowl and snapping turtles due to PCBs.

Services these natural resources provide to humans have been lost. For example, recreational fishing has been impaired by restrictions on taking fish from certain areas of the Hudson River. Further, PCB contamination in the Hudson River is a potential health threat to people who eat fish or who eat other organisms that inhabit the river and the surrounding area.

Living resources at every level of the Hudson's aquatic, terrestrial, and wetland-based food chains are contaminated with PCBs. PCB contamination is found in invertebrates, amphibians, reptiles, birds, and mammals such as mink, otter, bats, mice, shrews, and voles. PCB concentrations in wildlife exceed effects thresholds from the scientific literature. Such exceedances include the following:

- In sediments, PCBs are present at levels potentially causing harmful impacts to aquatic ecosystems.
- In fish, PCB levels associated with biochemical changes and adverse reproductive effects are exceeded.
- In mink, PCB levels associated with reproductive impairment are exceeded.
- In snapping turtles, PCB levels associated with the latent mortality in juveniles are exceeded.
- In bullfrogs, PCB levels associated with significant risk for various adverse effects in amphibians, including physical malformations, are exceeded.
- In birds, PCB levels associated with reproductive impairment are exceeded.

Serious adverse effects are likely to be occurring to these, and potentially other, living organisms exposed to the PCB contamination in the Hudson River region. To elucidate those effects, further studies on natural resources of the Hudson River, including fish, mink, sediment, birds, and waterfowl, are currently underway.

In conclusion, PCBs released from GE facilities on the Upper Hudson River present a serious and long-term threat to the health of the entire Hudson River ecosystem that *warrants continued study and further action to restore and compensate* for the injured natural resources and the services that have been lost. Because of concerns about the contamination and its potential impact, the Hudson River Natural Resource Trustees are continuing to assess the Hudson River ecosystem. The Trustees will use the information they collect during this assessment to document injuries to natural resources and determine the amount and type of restoration needed to compensate the public for these injuries.

FURTHER INFORMATION

Further information on the Hudson River natural resource damage assessment (NRDA) can be found at the following websites:

<http://www.darp.noaa.gov/northeast/udson/index.html>

<http://www.dec.ny.gov/lands/25609.html>

<http://www.fws.gov/contaminants/restorationplans/HudsonRiver/index.html>

To add yourself to the Hudson-NRDA listserv:

1. Send a message to: requests@willamette.nos.noaa.gov
2. Write in the subject: Subscribe hudsonnrda

If you have questions about natural resource damages, please contact one of the individuals listed below:

Tom Brosnan
National Oceanic and Atmospheric Administration
1305 East West Highway SSMC4, Room 10219
Silver Spring, MD 20910
301-713-3038 x186
Tom.Brosnan@noaa.gov

Kathryn Jahn
United States Fish and Wildlife Service
3817 Luker Road
Cortland, NY 13045
607-753-9334
Kathryn_Jahn@fws.gov

Sean Madden
New York State Department of Environmental Conservation
Division of Fish, Wildlife, and Marine Resources
625 Broadway, 5th Floor
Albany, NY 12233
518-402-8977
ssmadden@gw.dec.state.ny.us

REFERENCES

- Agency for Toxic Substances and Disease Registry (ATSDR). 2000. Public health statement: Polychlorinated biphenyls (PCBs). November.
- Bleavins, M.R., R.J. Aulerich, and R.K. Ringer. 1981. Placental and mammary transfer of polychlorinated and polybrominated biphenyls in the mink and ferret. *In*: Lamb D.W., Kenaga E.E., eds., *Avian and Mammalian Wildlife Toxicology*. American Society for Testing and Materials, Philadelphia, PA, pp. 121–131.
- Beckett, K.J., S.D. Millsap, A.L. Blankenship, M.J. Zwernick, J.P. Giesy, and S.J. Bursian. 2005. Squamous epithelial lesion of the mandibles and maxillae of wild mink (*Mustela vison*) naturally exposed to polychlorinated biphenyls. *Environ. Toxicol. Chem.* 24:674-677.
- Bopp, R.F., H.J. Simpson, and B.L. Deck. 1985. Final Report NYS C00708. Release of polychlorinated biphenyls from contaminated Hudson River sediments. Prepared for New York State Department of Environmental Conservation.
- Bursian, S.J., C. Sharma, R.J. Aulerich, B. Yamini, R.R. Mitchell, K.J. Beckett, C.E. Orazio, D. Moore, S. Svirsky, and D.E. Tillitt. 2006. Dietary exposure of mink (*Mustela vison*) to fish from the Housatonic River, Berkshire County, Massachusetts, USA: Effects on organ weights and histology and hepatic concentrations of polychlorinated biphenyls and 2,3,7,8-tetrachlorodibenzo-*p*-dioxin toxic equivalence. *Environ. Toxicol. Chem.* 25 (6):1541-1550.
- Bursian, S.J., J. Kern, R.E. Remington, J.E. Link, and S.D. Fitzgerald. 2011. Dietary exposure of mink (*Mustela vison*) to fish from the Upper Hudson River, New York, USA. SETAC North America, 32nd Annual Meeting, 13-17 November, Boston, MA. Available at: <http://www.fws.gov/contaminants/restorationplans/HudsonRiver/docs/Mink_reproduction_platform_SETAC_2011.pdf>
- Cignoll, M. 2012, January 17. Officials: Price increase for consumers the best option. *The Troy Record*, p. 6. Also available at <<http://www.troyrecord.com/articles/2012/01/26/river/doc4f2155ef056e7670943742.txt>>.
- Custer, C.M., T. Custer, and P. Dummer. 2010a. Patterns of organic contaminants in eggs of an insectivorous, an omnivorous, and a piscivorous bird nesting on the Hudson River, New York, USA. *Environ. Toxicol. Chem.* 29(10):2286-2296.
- Custer, T., C. Custer, and B. Gray. 2010b. Polychlorinated biphenyls, dioxins, furans, and organochlorine pesticides in spotted sandpiper eggs from the Upper Hudson River Basin, New York. *Ecotoxicology.* 19:391-404.
- Custer, T., C. Custer and B. Gray. 2010c. Polychlorinated biphenyls, dioxins, furans, and organochlorine pesticides in belted kingfisher eggs from the Upper Hudson River Basin, New York, USA. *Environmental Toxicology and Chemistry.* 29(1):99-110.
- Efroymsen, R.A., G.W. Suter II, B.E. Sample, and D.S. Jones. 1997. Preliminary remediation goals for ecological endpoints. Oak Ridge National Laboratory. ES/ER/TM- 162/R2. <http://www.hsrdr.ornl.gov/ecorisk/guidance.html>.
- Eisenreich, K.M., S.M. Kelley, and C.L. Rowe. 2009. Latent mortality of juvenile snapping turtles from the Upper Hudson River, New

- York, exposed maternally and via the diet to polychlorinated biphenyls (PCBs). *Environ. Sci. Technol.* 43(15):6052-6057.
- EPA (U.S. Environmental Protection Agency). 2000a. Hudson River PCBs Reassessment RI/FS Phase 3 Report: Feasibility Study. Prepared by TAMS Consultants, Inc. December.
- EPA (U.S. Environmental Protection Agency). 2000b. Database for the Hudson River PCBs Reassessment RI/FS. Release 5.0. October. Prepared by TAMS Consultants, Inc., Bloomfield, NJ.
- EPA (U.S. Environmental Protection Agency). 2002. Record of Decision on Hudson River PCBs Site. Region 2. United States Environmental Protection Agency, New York, NY.
- Foley, R.E., S.J. Jackling, R.J. Sloan, and M.K. Brown. 1988. Organochlorine and mercury residues in wild mink and otter: Comparisons with fish. *Environ. Toxicol. Chem.* 7:363-374.
- Funicello, J. 1979, June 25. State warns against eating snapping turtles. *Albany Times Union*, pp. 3, 5.
- Harris, M.L. and J.E. Elliott. 2011. Effects of polychlorinated biphenyls, dibenzo-*p*-dioxins and dibenzofurans, and polybrominated diphenyl ethers in wild birds. Chapter 14 in *Environmental Contaminants in Biota*, W.N. Beyer and J.P. Meador (eds.). CRC Press, Boca Raton, FL. pp. 477-528.
- Heaton, S.N., S.J. Bursian, J.P. Giesy, D.E. Tillitt, J.A. Render, P.D. Jones, D.A. Vergrugge, T.J. Kubiak, and R.J. Aulerich. 1995. Dietary exposure of mink to carp from Saginaw Bay, Michigan. 1. Effects on reproduction and survival, and the potential risks to wild mink populations. *Arch. Environ. Contam. Toxicol.* 28:334-343.
- HRNRT (Hudson River Natural Resource Trustees). 2001. Sampling and analysis plan, Hudson River fish health assessment phase I: Field sampling, necropsy, histopathology, disease, fish age (field version). Hudson River Natural Resource Damage Assessment. Final. October 3. Viewed at <http://www.dec.ny.gov/docs/wildlife_pdf/hrfishsap_rep.pdf>.
- HRNRT (Hudson River Natural Resource Trustees). 2002. Preliminary investigation of snapping turtles. Hudson River Natural Resource Damage Assessment. Summer. Viewed at: <<http://www.fws.gov/contaminants/restorationplans/HudsonRiver/HudsonRiverTurtleFactsheet.pdf>>.
- HRNRT (Hudson River Natural Resource Trustees). 2004a. Work summary and data report for the collection of eggs from American peregrine falcon, Hudson River, New York. December 24. Hudson River Natural Resource Damage Assessment. Available at: <<http://www.fws.gov/contaminants/restorationplans/HudsonRiver/docs/HudsonRiverPFalcoEggSumDataFINAL.pdf>>.
- HRNRT (Hudson River Natural Resource Trustees). 2004b. Study plan for year 2004 avian investigations for the Hudson River. Hudson River Natural Resource Damage Assessment. Final. June 15. Available at <http://www.darrp.noaa.gov/northeast/hudson/pdf/04avian_investigations.pdf>.
- HRNRT (Hudson River Natural Resource Trustees). 2005a. Data report for the collection of eggs from the common snapping turtle (*Chelydra serpentina serpentina*) from the Hudson River, New York. Hudson River Natural Resource Damage Assessment. March 30. Available at: <<http://www.fws.gov/contaminants/restorationplans/HudsonRiver/docs/SnappingTurtleEggDataReport.pdf>>.

- HRNRT (Hudson River Natural Resource Trustees). 2005b. Data report for the collection of eggs from spotted sandpipers, American woodcock, belted kingfisher, American robin, red-winged blackbird, and eastern phoebe associated with the Hudson River from Hudson Falls to Schodack Island, New York. Hudson River Natural Resource Damage Assessment. June 15. Available at: <http://www.darrp.noaa.gov/northeast/udson/pdf/avian_egg_rev_final_june_15_2005.pdf>
- HRNRT (Hudson River Natural Resource Trustees). 2005c. Data report for the collection of eggs from eastern screech owl associated with the Hudson River from Hudson Falls to Schodack Island, New York. Hudson River Natural Resource Damage Assessment. April 28. Available at: <<http://www.fws.gov/contaminants/restorationplans/udsonriver/ScreechOwlEggDataReportFINAL.pdf>>
- HRNRT (Hudson River Natural Resource Trustees). 2005d. Modification to study plan for avian investigations for the Hudson River USGS study plan amendment for 2005. Hudson River Natural Resource Damage Assessment. May 4. Available at <http://www.darrp.noaa.gov/northeast/udson/pdf/04avian_investigations.pdf>.
- HRNRT (Hudson River Natural Resource Trustees). 2006a. Study plan for mink injury investigations for the Hudson River. Hudson River Natural Resource Damage Assessment. Available at: <<http://www.fws.gov/contaminants/restorationplans/udsonriver/StudyPlanMink.pdf>>.
- HRNRT (Hudson River Natural Resource Trustees). 2006b. Modification to study plan for avian investigations for the Hudson River. USGS study plan amendment for 2006. Hudson River Natural Resource Damage Assessment. Final. May 9. Available 1-14-11 at <http://www.darrp.noaa.gov/northeast/udson/pdf/modifications_avian_study_plan_2006.pdf>.
- HRNRT (Hudson River Natural Resource Trustees). 2007a. Avian injury study: Avian egg injection study plan. Amendment for year 2 (2007). Hudson River Natural Resource Damage Assessment. June 1. Available at: <http://www.darrp.noaa.gov/northeast/udson/pdf/Final_Study_Plan_Avian_Injury_Study2_2007.pdf>.
- HRNRT (Hudson River Natural Resource Trustees). 2007b. Data report for the collection of bullfrog (*Rana catesbeiana*) tadpoles and near-shore sediment samples from the Hudson River, New York. Hudson River Natural Resource Damage Assessment. May. Available at: <<http://www.fws.gov/contaminants/restorationplans/HudsonRiver/docs/HudsonBullfrogTadpoleandSedimentReportPRVFINAL.pdf>>.
- HRNRT (Hudson River Natural Resource Trustees). 2007c. Work summary and data report: Collection of bats from the Hudson River, New York 2001 and 2002 samples. Hudson River Natural Resource Damage Assessment. September. Available at: <<http://www.fws.gov/contaminants/restorationplans/HudsonRiver/docs/HudsonRiverBatReportFINALPRV.pdf>>.
- HRNRT (Hudson River Natural Resource Trustees). 2007d. Study plan for avian egg injection study. Hudson River Natural Resource Damage Assessment. Final. Public Release Version. Released May 12, 2006. Revised January 31, 2007. U.S. Department of Commerce, Silver Spring,

- MD. Available at < http://www.dec.ny.gov/docs/wildlife_pdf/wpavwldin.pdf>.
- HRNRT (Hudson River Natural Resource Trustees). 2008a. Data report for the preliminary investigation of amphibian breeding habitat and screening of breeding pool sediments for polychlorinated biphenyl contamination, Hudson River, New York. Available at <<http://www.fws.gov/contaminants/restorationplans/HudsonRiver/docs/2004AmphibianHabitatandSedimentsDataReportFINAL.pdf>>
- HRNRT (Hudson River Natural Resource Trustees). 2008b. Fact sheet: Sediment toxicity pilot study for the Hudson River NRDA. September. Available at: <<http://www.fws.gov/contaminants/restorationplans/HudsonRiver/docs/SedimentToxicityPilotStudyFactSheetFINALSeptember2008.pdf>>
- HRNRT (Hudson River Natural Resource Trustees). 2008c. Fact sheet: Amphibian Investigations for the Hudson River NRDA. June. Available at: <<http://www.fws.gov/contaminants/restorationplans/HudsonRiver/docs/AmphibianHabitatandSedimentFACTSHEETJune2008.pdf>>
- HRNRT (Hudson River Natural Resource Trustees). 2008d. Injury determination report: Hudson River surface water resources. New York State Department of Environmental Conservation, National Oceanic and Atmospheric Administration, and U.S. Department of the Interior. Issued as part of the Hudson River Natural Resource Damage Assessment.
- HRNRT (Hudson River Natural Resource Trustees). 2008e. Study plan for avian injury study. Year 3 (2008). Hudson River Natural Resource Damage Assessment. June 30. Available at: <<http://www.fws.gov/Contaminants/restorationplans/HudsonRiver/AvianInjuryStudyPlanFinalJune2008.pdf>>.
- HRNRT (Hudson River Natural Resource Trustees). 2008f. Study plan for waterfowl injury assessment: Determining PCBs concentrations in Hudson River resident waterfowl. Final. December. Available at: <<http://www.fws.gov/contaminants/restorationplans/HudsonRiver/docs/StudyPlanPCBsHudsonRvrResidentWaterfowlFINALDec08.pdf>>.
- HRNRT (Hudson River Natural Resource Trustees). 2009a. Data report: Organochlorine and metal contaminant levels in Hudson River aquatic insects. Hudson River Natural Resource Damage Assessment. September. Available at: <<http://www.fws.gov/contaminants/restorationplans/HudsonRiver/docs/aquaticinsectdr.pdf>>.
- HRNRT (Hudson River Natural Resource Trustees). 2009b. Fact sheet: Fish toxicity pilot study for the Hudson River NRDA. May. Available at: <<http://www.fws.gov/contaminants/restorationplans/HudsonRiver/docs/SturgeonToxicityTestingFactSheetFINALMay2009.pdf>>
- HRNRT (Hudson River Natural Resource Trustees). 2009c. Study plan for avian injury study year 4 (2009). Hudson River Natural Resource Damage Assessment. June 11. Available at: <http://www.fws.gov/contaminants/restorationplans/hudsonriver/docs/Hudson_River_Avian_Injury_Study_Year_4_Trustee_Study_Plan_FINAL.pdf>.
- HRNRT (Hudson River Natural Resource Trustees). 2010. Data report for the collection of small mammals and American woodcock from the floodplain of the Hudson River, New York in year 2001, analysis of floodplain earthworms from the

year 2000, and re-analysis of select floodplain soils and small mammals from the year 2000. Hudson River Natural Resource Damage Assessment. January. Available at: < http://www.dec.ny.gov/docs/wildlife_pdf/fp0100.pdf >.

- HRNRT (Hudson River Natural Resource Trustees). 2011a. Congener-specific analysis of polychlorinated biphenyl residues in tree swallow chicks, eggs and other biota from the Hudson River, W.U. 30096. Report dated November 25, 1996. Final Laboratory Report FY-97-30-01.
- HRNRT (Hudson River Natural Resource Trustees). 2011b. U.S. Geological Survey, July 2000, Organochlorine contaminants in biota from the Hudson River, New York. USGS Final Report FY-99-31-01, FWS No. 14-48-0005-50181-97-J-008, CERC No: 82070-1491CL62. FWS Project Title "Investigation of exposure of migratory birds to PCBs, PCDDs, PCDFs and organochlorine pesticides along the Hudson River." August 1, 2011. Final. U.S. Department of Commerce, Silver Spring, MD.
- HRNRT (Hudson River Natural Resource Trustees). 2011c. U.S. Geological Survey, October 2000, Organochlorine contaminants in bald eagle eggs. USGS Final Report FY-00-31-04, FWS No. 1448-50181-99-H-007, CERC No. 3307-70L1D. FWS Project Title: "Chemical contamination of nesting tree swallows, great blue herons, and resident/nesting bald eagles along the Hudson River, New York." Final. August 1, 2011. U.S. Department of Commerce, Silver Spring, MD.
- HRNRT (Hudson River Natural Resource Trustees). 2011d. Organochlorine contaminants in tree swallow nestlings and in adipose tissue from great blue heron nestlings. FY-00-31-05. FWS NO: 1448-50181-99-H-007. CERC NO: 3307-70L1D.

FWS Project Title: "Chemical Contamination of Nesting Tree Swallows, Great Blue Herons, and Resident/Nesting Bald Eagles Along the Hudson River, New York." Report dated November 2000.

- HRNRT (Hudson River Natural Resource Trustees). 2011e. U.S. Geological Survey, July 2003, Polychlorinated biphenyls and organochlorine pesticides in bald eagle blood and egg samples from the Hudson River, New York. USGS Report CERC-8335-FY03-31-04, FWS No. 50181-1-H014, DNC No. 50181-1-H014A. FWS Project Title: "Chemical contamination of resident/nesting bald eagles along the Hudson River, New York – Samples from 1999-2001." Includes correction. Final. August 1, 2011. U.S. Department of Commerce, Silver Spring, MD.
- HRNRT (Hudson River Natural Resource Trustees). 2011f. U.S. Geological Survey, February 2003, Polychlorinated biphenyls and organochlorine pesticides in bald eagles and fish from the Hudson River, New York, sampled 1999-2001. USGS Report CERC-8335-FY03-31-02, FWS No. 50181-1-H014, DNC No. 50181-1-H014A. FWS Project Title: "Chemical contamination of resident/nesting bald eagles along the Hudson River, New York – Samples from 1999-2001." Final. August 1, 2011. U.S. Department of Commerce, Silver Spring, MD.
- HRNRT (Hudson River Natural Resource Trustees). 2011g. U.S. Geological Survey, September 2005, Polychlorinated biphenyls, organochlorine pesticides, dioxins and furans in bald eagle egg and blood samples from the Hudson River, New York. USGS Report CERC-8335-FY05-31-03. Final. August 1, 2011. U.S. Department of Commerce, Silver Spring, MD.
- HRNRT (Hudson River Natural Resource Trustees). 2011h. Fact sheet: Mink injury investigations for the Hudson River NRDA

- 2012-2013. March. Available at <<http://www.fws.gov/contaminants/restorationplans/HudsonRiver/docs/MinkFactSheetMarch2012.pdf>>.
- HRNRT (Hudson River Natural Resource Trustees). 2011i. Study plan for mink injury determination. Investigation of mink abundance and density relative to polychlorinated biphenyl contamination within the Hudson River. Hudson River Natural Resource Damage Assessment. March 19. Draft for Public Review and Comment. Available at: <http://www.fws.gov/contaminants/restorationplans/HudsonRiver/docs/TrusteeStudyPlanMinkFieldStudy_03192012.pdf>.
- HRTC (Hudson River Trustee Council). 2002. Hudson River Natural Resource Damage Assessment Plan. September.
- Hugla, J.L. and J.P. Thomé. 1999. Effects of polychlorinated biphenyls on liver ultrastructure, hepatic monooxygenases, and reproductive success in the barbell. *Ecotoxicology and Environmental Safety* 42(3):265-273.
- Kelly, S.M., K.M. Eisenreich, J.E. Baker, and C.L. Rowe. 2008. Accumulation and maternal transfer of polychlorinated biphenyls in snapping turtles of the Upper Hudson River, New York, USA. *Environ. Toxicol. Chem.* 27(12):2565-2574.
- Kim K.S., Pastel M.J., Kim J.S., and Stone W.B. 1984. Levels of polychlorinated biphenyls, DDE, and mirex in waterfowl collected in New York State, 1979-1980. *Arch Environ Contam Toxicol* 13:373-381.
- Kim, H.T., K.S. Kim, J.S. Kim, and W.B. Stone. 1985. Levels of polychlorinated biphenyls, DDE, and mirex in waterfowl collected in New York State, 1981-1982. *Arch. Environ. Contam. Tox.* 14:13-18.
- Litten, S. 2003. Contaminant Assessment and Reduction Project Water (CARP). New York State Department of Environmental Conservation. Viewed at <<http://www.dec.ny.gov/chemical/23839.html>>.
- Malcolm Pirnie, Inc. 2009. Village of Stillwater well field investigation report. Draft final. U.S. Army Corps of Engineers – Kansas City District. U.S. Environmental Protection Agency – Region 2. February.
- Mayack, D.T. and J. Loukmas 2001. Progress report on Hudson River mammals: Polychlorinated biphenyl (PCB) levels in mink, otter, and muskrat and trapping results for mink, the upper Hudson River drainage, 1998-2000. March.
- McCarty, J.P. and A.L. Secord. 1999a. Reproductive ecology of tree swallows (*Tachycineta bicolor*) with high levels of polychlorinated biphenyl contamination. *Environ. Toxicol. Chem.* 18(7):1433-1439.
- McCarty L.P. and A.L. Secord. 1999b. Nest-building behavior in PCB contaminated tree swallows. *Auk* 116:55–63.
- McCarty J.P. and A.L. Secord. 2000. Possible effects of PCB contamination on female plumage color and reproductive success in Hudson River tree swallows. *The Auk* 117 (4):987-995.
- Niimi, A.J. 1996. PCBs in aquatic organisms. In: *Environmental Contaminants in Wildlife: Interpreting Tissue Concentrations*. W.N. Beyer, G.H. Heinz, and A.W. Redmon-Norwood (eds.). CRC Press, Boca Raton, FL. p. 117-152.
- NOAA (National Oceanic and Atmospheric Administration). 2010. Hudson River Query Manager database, updated January 15, 2010. Latest version available at: <<http://response.restoration.noaa.gov/watersheddownloads>>.
- NYSDEC (New York State Department of Environmental Conservation). 1999. Technical guidance for screening contaminated sediments. Division of Fish

- and Wildlife, Division of Marine Resources. Sloan, R.J., M.W. Kane and L. Skinner. 2005. Of time, PCBs, and the fish of the Hudson River. Bureau of Habitat, Division of Fish, Wildlife and Marine Resources, New York State Department of Environmental Conservation, Albany, New York. http://www.dec.ny.gov/docs/wildlife_pdf/hrpcbtrend.pdf
- NYSDEC (New York State Department of Environmental Conservation). 2004. Record of Decision. GE Hudson Falls Plant Site, Operable Units No. 2A-2D. Village of Hudson Falls, Town of Kingsbury, Washington County, New York. Site Number 5-58-013. March.
- NYSDOH (New York State Department of Health). 1993. Health Advisories. Chemicals in sportfish and game: 1993-1994.
- NYSDOH (New York State Department of Health). 2008. Chemicals in sportfish and game: 2008-2009 health advisories.
- NYSDOH (New York State Department of Health). 2011. Chemicals in sportfish and game: 2011-2012 health advisories.
- Restum, J.C., S.J. Bursian, J.P. Giesy, J.A. Render, W.G. Helferich, E.B. Shipp, and D.A. Verbrugge. 1998. Multigenerational study of the effects of consumption of PCB-contaminated carp from Saginaw Bay, Lake Huron, on mink. 1. Effects on mink reproduction, kit growth and survival, and selected biological parameters. *J. Toxicol. Environ. Health, Part A* 54:343-375.
- SEA Consultants, Inc. 2002. Hudson River natural resource damage assessment floodplain soil and biota screening sampling report. Prepared for Industrial Economics, Inc. under contract to the National Oceanic and Atmospheric Administration. February.
- Secord, A.L. and J.L. McCarty. 1997. Polychlorinated biphenyl contamination of tree swallows in the Upper Hudson River Valley, New York. Effects on breeding biology and implications for other bird species. New York Field Office, U.S. Fish and Wildlife Service, Cortland, NY.
- Smit, M.D., P.E.G. Leonards, A.J. Murik, A.W.J.J. de Jongh, and B. van Hattum. 1996. Development of otter-based quality objectives for PCBs, Institute for Environmental Studies, Vrije Universiteit, Amsterdam.
- Weston Solutions, Inc. 2003. Ecological risk assessment for General Electric (GE)/Housatonic River Site, Rest of River. Appendix E: Assessment endpoint – Community condition, survival, reproduction, development, and maturation of amphibians. Prepared for the U.S. Army Corps of Engineers and U.S. Environmental Protection Agency. July.
- Weston Solutions, Inc. 2005. Floodplain soil sampling summary report, Hudson River PCBs Site, New York. Prepared for the U.S Environmental Protection Agency. W.O. No. 20103.001.001.1048.00. August.



MODIFICATION TO STUDY PLAN FOR MINK INJURY DETERMINATION

INVESTIGATION OF MINK ABUNDANCE AND DENSITY RELATIVE TO POLYCHLORINATED BIPHENYL CONTAMINATION WITHIN THE HUDSON RIVER DRAINAGE

HUDSON RIVER NATURAL RESOURCE DAMAGE ASSESSMENT

HUDSON RIVER NATURAL RESOURCE TRUSTEES

STATE OF NEW YORK

U.S. DEPARTMENT OF COMMERCE

U.S. DEPARTMENT OF THE INTERIOR

AUGUST 15, 2013

Available from:
U.S. Department of Commerce
National Oceanic and Atmospheric Administration
Hudson River NRDA, Lead Administrative Trustee
Damage Assessment Center, N/ORR31
1305 East-West Highway, Rm 10219
Silver Spring, MD 20910-3281



**MODIFICATION TO STUDY PLAN FOR MINK INJURY DETERMINATION
INVESTIGATION OF MINK ABUNDANCE AND DENSITY RELATIVE TO POLYCHLORINATED BIPHENYL
CONTAMINATION WITHIN THE HUDSON RIVER DRAINAGE**

INTRODUCTION

Past and continuing discharges of polychlorinated biphenyls (PCBs) have contaminated the natural resources of the Hudson River. The Hudson River Natural Resource Trustees – New York State, the U.S. Department of Commerce, and the U.S. Department of the Interior – are conducting a natural resource damage assessment (NRDA) to assess and restore those natural resources injured by PCBs (Hudson River Natural Resource Trustees 2002).

Many species of mammals rely on the Hudson River, including its floodplain, for habitat, food, and as a breeding ground. Mammals that depend on the river for food and habitat include otter, muskrat, raccoon, beaver, and mink. The Hudson River NRDA Plan identified mink health as an area of biological injury investigation.

On August, 2, 2010, the Trustees released a Draft Study Plan entitled, “Investigation of Mink Occupancy Relative to Polychlorinated Biphenyl Contamination within the Hudson River Drainage” (Hudson River Natural Resource Trustees, 2010). Following peer and public review of that plan, the Trustees determined that revisions to that plan were appropriate, resulting in the March 19, 2012 Draft Study Plan (Hudson River Natural Resource Trustees, 2012a) being released for further peer and public review, culminating in a Final Study Plan released on July 13, 2012 (Hudson River Natural Resource Trustees, 2012b).

As outlined in the Final Study Plan, the summer of 2012 served as a pilot study to inform the design of the 2013 sampling season. The modifications below pertain to 2013.

The Trustees have evaluated the changes to the study described in this Study Plan Modification for 2013 and determined that the changes are not sufficiently substantive to necessitate peer and public review of the Study Plan Modification for Year 2013.

MODIFICATIONS

Appendix A - STUDY PLAN INVESTIGATION OF MINK ABUNDANCE RELATIVE TO POLYCHLORINATED BIPHENYL (PCB) CONTAMINATION WITHIN THE HUDSON RIVER DRAINAGE

Page 10: The number of sample sites the scat dogs will survey has been changed from 144 sites in each area to 75 sites in the Hudson drainage and 70 sites in the Mohawk drainage. Any other references in the Final Study Plan to the number of sample sites for scat surveys will be changed from 144 sites in each area to 75 sites in the Hudson drainage and 70 sites in the Mohawk drainage. The number of sample sites for hair snares has been changed from 50 sites per river to 71 sites in the Hudson drainage and 47 sites in the Mohawk drainage.

Page 12: The study areas will no longer be divided into standard 1 km x 1 km grid cells with one potential sample location per grid cell. Instead, samples will be clustered.

Page 12: The hair snare design referenced in Appendix 1 was further refined after the 2012 field season. The changes in dimensions are listed under the modifications to Appendix 7. Sardines and mink gland lure will no longer be used in the hair snares; instead the traps will be baited with drops of a fish oil mink bait.

Page 13: If hairs are seen on a gun brush, the whole gun brush will no longer be removed. Instead, technicians will use tweezers to remove hairs from the gun brush, and hairs will be placed into labeled coin envelopes. The labeled coin envelopes will be sealed and placed with a desiccant in a plastic bag. After removing any hairs from a gun brush the technicians will hold the gun brush and tweezers to a flame (such as from a lighter) for 30 seconds to remove any stray debris, allowing continued use of the equipment.

If there are fewer than 2 hairs total in a single hair snare, the hairs will not be collected. If there are 2 or more hairs in a brush, the hairs will be collected, processed, and sent to the lab in the same envelope. If there are 2 or more hairs from a combination of the brushes and the sides of the trap they will be collected, processed and sent to the lab in different envelopes representing where they were found (brush 1, brush 2, side of trap).

Page 14: When scats are collected in the field they will no longer be placed in waxless bags and dried for 1-4 days. When scats are collected in the field, they will immediately be placed in falcon tubes of ethanol. Scat will be stored in 95% ethanol, not 96% ethanol.

Appendix 5, page 56: The collection of habitat variables during scat collection will be simplified. These changes will also be reflected in the data sheets.

Appendix 7, page 67: Hair snares will be both anchored and camouflaged with natural materials (logs, rocks, etc.), not anchored using tent stakes or water bottles. For the 2013 sampling season the hair snare dimensions will be approximately 80 cm x 45 cm, folded every 15 cm to make a triangular opening. The position of the gun brushes will be 30 cm in from the openings. A fish oil bait will be used in 2013. Hair snares will be visited every 14 days, not every 7 days as originally planned.

Appendix 8, page 70: The hair collection data sheet has been revised. Appendix A contains a copy of the new hair collection data sheet.

Appendix 10, page 72: As above, scats will no longer be placed in waxless bags and dried. Scats will be placed in ethanol in the field.

Appendix 11, page, 77: As above, scats will no longer be placed in waxless bags and dried. Scats will be placed in ethanol in the field.

Appendix 12, page 79: Mink scat data sheets will be changed to include the simplified collection of habitat variables. Appendix A contains a copy of the new scat collection data sheet.

APPENDIX A

REVISED HAIR COLLECTION DATA SHEET

REVISED SCAT COLLECTION DATA SHEET

**MODIFICATION TO STUDY PLAN FOR MINK INJURY DETERMINATION
INVESTIGATION OF MINK ABUNDANCE AND DENSITY RELATIVE TO POLYCHLORINATED BIPHENYL
CONTAMINATION WITHIN THE HUDSON RIVER DRAINAGE**

2013 HR Mink Abundance Study - Scat Collection

Site Info		Weather		Habitat Observations									
Date:		Temp (°C):		Upstream				Downstream					
Site ID:		Wind Speed (m/s):		Stream Section		Ease of access	Bank slope index	Stream Width	Avg Water Depth	Ease of access	Bank slope index	Stream Width	Avg Water Depth
Recorder:		Precip:		0 - 50m									
Dog Team:		None Humid Drizzle Rain		50 - 100m									
Scat Collector:		Cloud Cover (%):		100 - 150m									
Start time:		General Observations		150 - 200m									
Waypoint (road):				200 - 250m									
WPt Start (road):				250 - 300m									
WPt End (up):				300 - 350m									
WPt Start (road):				350 - 400m									
WPt End (down):						E = Easy M = Medium D = Difficult	F = Flat, 5°-45°, 45°-85°, V = Vertical	0-5ft, 5-10ft, 10-15ft,etc...	0-1ft, 1-2ft,2-3ft, 3-4ft,4-5ft, >5ft	E = Easy M = Medium D = Difficult	F = Flat, 5°-45°, 45°-85°, V = Vertical	0-5ft, 5-10ft, 10-15ft,etc...	0-1ft, 1-2ft,2-3ft, 3-4ft,4-5ft, >5ft
End time:													

Scat Collection

Time	Scat Sample	GPS # (waypoint)	Photo	Degree of Confidence	Description of scat location	Dist. to Water (m)	Scat Size (mm)		Scat Condition		Reward given to dog?	Notes
							Length	Width	Freshness	Color		
hh:mm	e.g. A,B,...,ZZ		e.g. DSCN0309	H = High M = Medium L = Low	leaf litter; needle litter; log; rock; latrine; brush; buried				M= moist/fresh DS = dry ext. moist/Soft int DF = dry ext. moist/Firm int D = dry throughout (crumbly)	Bl = Black GBI = Green/Black G = Green Br = Brown Gy = Grey Other	Yes/No	

Scat Collection

Time	Scat Sample	GPS # (waypoint)	Photo	Degree of Confidence	Description of scat location	Dist. to Water (m)	Scat Size (mm)		Scat Condition		Reward given to dog?	Notes
							Length	Width	Freshness	Color		
hh:mm	e.g. A.B....ZZ		e.g. DSCN0309	H = High M = Medium L = Low	leaf litter; needle litter; log; rock; latrine; brush; buried				M = moist/fresh DS = dry ext. moist/Soft int DF = dry ext. moist/Firm int D = dry throughout (crumbly)	Bl = Black GBL = Green/Black G = Green Br = Brown Gy = Grey Other	Yes/No	



FACT SHEET

HUDSON RIVER

RESTORATION PLANNING



HUDSON RIVER TRUSTEES

ASSESSING AND RESTORING YOUR NATURAL RESOURCES

Publication of List of Restoration Project Proposals Submitted by Public September 2013

Restoration is the goal of the Hudson River Natural Resource Damage Assessment. Restoration is an active component of damage assessment that can be seen and felt for generations.

The Hudson River Trustees received a request for a list of restoration projects that have been submitted by the public. The list (Table 1) released by the Trustees contains all of the projects received to date that have been submitted through the restoration project proposal form. For this list, please read Table 1: Hudson River Restoration Project Proposals from the Public; Table 1 is organized by county and individual projects have not been ranked or prioritized. Note that individual members of the public who submitted proposals are not named due to privacy concerns. The superscript letter (Member of the Public^a) can be used to track multiple projects received from the same individual.

Public meetings have given valuable input to the Trustees and have led to informal suggestions for restoration projects. Projects suggested at a public meeting are included in a separate list because an official restoration project proposal does not exist for such projects. For this list, please read Table 2: Hudson River Informal Restoration Project Proposals.

Projects on these lists will be considered for inclusion in the restoration plans the Trustees will develop. The Trustees are still actively accepting new restoration project proposal forms and encourage members of the public to submit new or updated restoration project proposals using the form available at www.darrp.noaa.gov/northeast/hudson/restore.html

Criteria for Restoration Project Assessment

During the damage assessment, Trustees will review the list of potential restoration projects, assessing the projects using criteria including, but not limited to:

- **Relevance-** Is there a sufficient connection between the proposed restoration effort and natural resource injuries and losses to the public?
- **Legality-** Does the restoration project comply with applicable/relevant federal, state, and local laws and regulations?
- **Efficacy-** How likely is it that the restoration project will be successful?
- **Cost Reasonableness-** What is the cost of the proposed restoration effort? Can the benefits be quantified? Is there an opportunity to share costs with other organizations and/or agencies?
- **Ecological Leverage-** Will the restoration project promote other environmental benefits?
- **Existing Plans-** Does the restoration project address or build upon existing federal, state, or local management plans?





HUDSON RIVER TRUSTEES

ASSESSING AND RESTORING YOUR NATURAL RESOURCES

Timing of Project Selection

The Trustees have not established a schedule for selecting restoration projects because of uncertainties surrounding the time needed to complete the injury analysis, evaluate restoration options and projects against the injuries, and resolve the damage claim. Our goal remains to begin restoration as soon as possible after the damage claim is resolved. Restoration projects will compensate the public for the loss of the Hudson River's natural resources and services from PCB contamination and remediation.

The Trustees will propose a list of potential restoration projects that will offset the public's loss, and those projects will be incorporated into a Draft Restoration Plan. The Trustees anticipate sharing this plan with you, the public, in meetings or other presentations, and soliciting your views on this information.

We are extremely grateful for the value that an informed and involved public brings to the process and to our decision-making.

More Information

www.fws.gov/contaminants/restorationplans/HudsonRiver/index/html

www.dec.ny.gov/lands/25609.html

www.darrp.noaa.gov/northeast/udson

To add yourself to the Hudson NRDA listserv:

1. Send a message to: requests@willamette.nos.noaa.gov
2. Write in subject: Subscribe hudsonnrda

If you have questions about natural resource damages, contact:

Tom Brosnan (NOAA), 301-713-3038 x186; Tom.Brosnan@noaa.gov

Kathryn Jahn (USFWS), 607-753-9334; Kathryn_Jahn@fws.gov

Sean Madden (NYSDEC), 518-402-8977; ssmadden@gw.dec.state.ny.us




Table 1: Hudson River Restoration Project Proposals from the Public

County	Town or Municipality	Project #	Project Description	Submitted By
Albany	Menands	A1	Make changes to the Route 387 Bridge, and Interstate 787 expressway and its ramps, to improve Hudson River access for pedestrians, cyclists, and people with disabilities.	Village of Menands
Albany	Menands	A2	Remove culverts to improve water flow between Little River and the estuary.	Village of Menands
Albany	Troy	A3	Restore Hudson River shoreline for swimming, boating, and fishing in Troy, NY.	City of Troy
Albany	Troy	A4	Clean up tributaries (Poestenkill, Wynatskill) and allow public access for fishing of migratory species like shad and herring.	City of Troy
Columbia	Hudson	C1	Create an environmental education center in Hudson, NY.	Columbia County EMC
Columbia	Stockport	C2	Dredge to deepen boat launch at the end of Station Road to allow access for fire protection.	Stockport Fire Co.
Dutchess	Fishkill	D1	Remove debris along Metro-North rails and develop greenway trail to improve access to Hudson River.	Town of Fishkill
Dutchess	Hyde Park	D2	Identify and remove flow and fishery impediments in Hudson River tributary Crum Elbow Creek, to improve habitat for fish and increase recreational fishing.	New York Rivers United
Dutchess	Location not specified.	D3	Remove water chestnut, an invasive species, from nontidal habitats.	Dutchess County Lakes Committee
Dutchess	Poughkeepsie	D4	Provide space (one square yard) for applicant to place non-denominational shrine to honor the Hudson River.	Member of the Public ^j
Dutchess	Rhinebeck	D5	Restore and remediate the Landsman Kill to improve the Hudson River watershed.	Member of the Public ^f
Dutchess	Location not specified.	D6	Support local stream watershed groups with funding for coordinator positions as well as educational materials, supplies, and monitoring equipment.	Dutchess County Environmental Management Council
Dutchess	Poughkeepsie	D7	Remove invasive water chestnut and sediment from Vanderburgh Cove to restore to historical conditions.	Vanderburgh Cove Neighborhood Association
Greene	Athens	G1	Restore natural stream channel pattern to improve water flow to wetland in Athens, NY.	Greene County Soil and Water Conservation District

Table 1: Hudson River Restoration Project Proposals from the Public

County	Town or Municipality	Project #	Project Description	Submitted By
Greene	Athens	G2	Stabilize banks and develop riparian buffers to decrease sedimentation in Catskill Creek and Kaaterskill Creek.	Greene County Soil and Water Conservation District
Greene	Athens	G3	Implement erosion control measures to slow migration of sediments from Sleepy Hollow Lake into Murderers Creek and the Hudson River.	Greene County Soil and Water Conservation District
Greene	Coxsackie	G4	Stabilize bank, improve boat access, and place informational kiosk on Coxsackie Island Preserve.	Greene County Soil and Water Conservation District
Greene	Coxsackie	G5	Improve access for canoes and kayaks in Coxsackie Creek.	Greene County Soil and Water Conservation District
Greene	New Baltimore	G6	Develop nature trails and environmental kiosks on Hudson River bank near New Baltimore, NY.	Greene County Soil and Water Conservation District
Greene	Athens/ Coxsackie	G7	Fix dyke to raise water level, improve habitat, and restore hydraulic function in Vosburgh Swamp.	Greene County Soil and Water Conservation District
Greene	Catskill	G8	Provide environmental education through interpretative signage at Cohotate Preserve.	Greene County Soil and Water Conservation District
Manhattan	New York City	M1	Improve waterfront access and increase water-based recreational opportunities in the northern portion of Riverside Park between 145th and 155th Streets. This will make the park more inviting and safe, and strengthen linkages within the residential and commercial communities near the park.	West Harlem Art Fund
Manhattan	New York City	M2	Implement Master Plan for West Harlem Waterfront Park and Piers in Manhattanville. Improve city-owned land on waterfront (125th to 135th Street between Broadway and the Hudson River), leading to environmental improvements and economic development of neighborhood.	West Harlem Art Fund
Orange	Beacon	O1	Restore 5 acres of degraded tidal wetlands on waterfront property owned by Scenic Hudson, to improve habitat and protect water quality. Surrounding area will be used for a public park and private development.	Scenic Hudson
Putnam	Cold Spring	P1	Purchase kayak to improve trash removal efforts on Hudson River.	Sons of the American Revolution, West/Putnam Chapter

Table 1: Hudson River Restoration Project Proposals from the Public

County	Town or Municipality	Project #	Project Description	Submitted By
Rensselaer	Rensselaer	R1	Develop wildlife viewing areas for falcon nesting at the Riverfront Park.	Member of the Public ^a
Rensselaer	Troy	R2	Identify and remove flow and fishery impediments in Hudson River tributary Wynants Kill, to improve habitat for fish and increase recreational fishing.	New York Rivers United
Rensselaer	Rensselaer	R3	Add ten acres to Papscaenee Island Nature Preserve, providing access to Hudson River for people with disabilities.	Rensselaer County Environmental Management Council
Rensselaer	Troy	R4	Identify and remove flow and fishery impediments in Hudson River tributary Poesten Kill, to improve habitat for fish and increase recreational fishing.	New York Rivers United
Rockland	Haverstraw	R5	Restore 600 linear feet of steel bulkhead, restore two inlets for public access, and create a doublewide boat launch to allow public access and enjoyment of river.	Rockland County Environmental Resources
Rockland	Nyack	R6	Establish a demonstration oyster bed restoration project, including local schools and community groups. Work toward the goal of eventually restoring this important fishery at Nyack Beach State Park and Memorial Park.	J.C. Brotherhood
Saratoga	Fort Edward	S1	Create parks for public access to capped sites at Fort Edward.	Environmental Advocates
Saratoga	Fort Edward	S2	Assess and dredge the Fort Edward Basin to reduce PCB exposure.	Member of the Public ^h
Saratoga	Location not specified.	S3	Implement projects listed in the town waterfront management plans.	Saratoga County Environmental Management Council
Saratoga	Mechanicville	S4	Construct gazebo near Champlain Canal wall to give community a platform for events and increase tourism.	Mechanicville Department of Public Works
Saratoga	Moreau	S5	Clean up PCB dredge spoil and develop into a park or flood plain area.	Member of the Public ^g
Saratoga	Saratoga	S6	Restore navigation to the historic Champlain Canal.	Town of Saratoga

Table 1: Hudson River Restoration Project Proposals from the Public

County	Town or Municipality	Project #	Project Description	Submitted By
Saratoga	Saratoga/ Northumberland/ Greenwich	S7	Construct Bi-County educational park focusing on ecological, geologic, and historic values of the Hudson River.	Saratoga County Chamber of Commerce
Saratoga	Schuylerville	S8	Restore contaminated river bottom and shoreline at the Schuylerville municipal beach.	Member of the Public ⁱ
Saratoga	Schuylerville	S9	Restore old Champlain Canal, harbor, and junction lock for use by recreational boaters and historical interpretation.	Member of the Public ⁱ
Saratoga	Schuylerville	S10	Construct environmental, recreational, and historical educational park to provide indoor and outdoor educational and research opportunities. Create an area where the public can go under the water level of Hudson River to observe natural processes.	Hudson Crossing Park Steering Committee
Saratoga	Schuylerville	S11	Restore contaminated river bottom and shoreline for the Schuylerville municipal beach.	Schuylerville Area Chamber
Saratoga	Schuylerville	S12	Remove contaminated sediments to re-open the Schuylerville Harbor and the canal between the harbor and the Hudson/Champlain Canal.	Village of Schuylerville
Saratoga	Schuylerville	S13	Extend the existing boardwalk on Schuylerville Municipal Beach 700 feet to reach the boat launch. Build a gazebo for public enjoyment and increase lighting for safety.	Village of Schuylerville
Saratoga	Schuylerville	S14	Restore contaminated river bottom and shoreline at the Schuylerville municipal beach.	Village of Schuylerville
Saratoga	Stillwater	S15	Rebuild seawall, boat launch and parking area.	Member of the Public ^c
Ulster	Esopus	U1	Identify and remove flow and fishery impediments in Hudson River tributary Black Creek, to improve habitat for fish and increase recreational fishing.	New York Rivers United
Ulster	Location not specified.	U2	Identify and remove flow and fishery impediments in Hudson River tributary Roundout Creek, to improve habitat for fish and increase recreational fishing.	New York Rivers United
Ulster and Orange	Location not specified.	U3	Identify and remove flow and fishery impediments in Hudson River tributary Quassaic Creek, to improve habitat for fish and increase recreational fishing.	New York Rivers United

Table 1: Hudson River Restoration Project Proposals from the Public

County	Town or Municipality	Project #	Project Description	Submitted By
Washington	Fort Edward	W1	Provide residents of Fort Edward with free municipal waste disposal at the Bi-County Incinerator in Hudson Falls. Recover the costs associated with the premature closing of solid waste landfills in Fort Edward.	Hudson River CARE
Washington	Fort Edward	W2	Provide residents and small businesses of Fort Edward with free municipal water. Reimburse town for costs of reestablishing the water district.	Hudson River CARE
Washington	Fort Edward	W3	Create an interpretive nature area and river front access park with restrooms, outdoor lighting, public boat launch, fishing pier, benches, signage, and parking lots. Improve public access roads and create hard surface parking area with lighting. Provide funds for the operation and upkeep of the park.	Hudson River CARE
Washington	Fort Edward	W4	Construct water treatment plant and air stripper to remove TCE from drinking water and provide safe water to the community.	Village of Fort Edward Board of Trustees
Washington	Fort Edward	W5	Replace existing municipal water lines, including a 20-inch water main (encased in cement at the bottom of the Hudson River) that is currently inaccessible for maintenance of repair, and a 6-inch feeder line to Rogers Island that is subject to frequent damage from boat traffic.	Village of Fort Edward Board of Trustees
Washington	Fort Edward	W6	Create a water/recreation park on the Northern edge of Rogers Island to provide enhanced public access to the Hudson River.	Village of Fort Edward Board of Trustees
Washington	Fort Edward	W7	Improve Fort Edward Yacht basin as a public arts and recreation park. Construct an amphitheater, provide picnic tables and benches, improve lighting, and provide sanitary services such as restrooms, showers, and a pump out station.	Village of Fort Edward Board of Trustees
Washington	Fort Edward	W8	Recover the expenses incurred for the design and construction of a \$5.2 million water treatment plant and the interest on the bond.	Hudson River CARE

Table 1: Hudson River Restoration Project Proposals from the Public

County	Town or Municipality	Project #	Project Description	Submitted By
Washington	Fort Edward	W9	Improve navigational dredging near the Route 197 Bridge to allow larger vessels to access the Fort Edward Yacht Basin.	Hudson River CARE
Washington	Fort Edward	W10	Improve the Fort Edward Yacht Basin for water-based recreation, including new docking stations and new pedestrian bridge to Rogers Island.	Hudson River CARE
Washington	Fort Edward	W11	Enhance Bradley Beach and Park on the North end of Rogers Island, including fishing pier, boat launch, and hard surfaced parking lot.	Hudson River CARE
Washington	Fort Edward	W12	Upgrade and maintain water lines in Fort Edward, including an enlarged water line and new pumping station and sewer line to Rogers Island, and replacement of 20-inch water line on the river bottom.	Hudson River CARE
Washington	Fort Edward	W13	Replace the existing PCB/TCE contaminated sewer lines from Defiance Asphalt, reducing public exposure to harm. Expand Fort Edward nature to join region's Feeder Canal trail system.	Hudson River CARE
Washington	Fort Edward	W14	Create a water containment system for McIntyre Park by diverting polluted spring away from the center of the park, making this a safer family recreation area. Install improved lighting, bleachers, and a hard-surfaced parking area.	Hudson River CARE
Washington	Fort Edward	W15	Create a recreation area on Bond Creek, including small docks for kayaks and canoes, nature/walking trails, and benches.	Hudson River CARE
Washington	Fort Edward	W16	Create a recreation area on Moses Creek, including small docks for kayaks and canoes, small fishing piers, nature/walking trails, and benches.	Hudson River CARE
Washington	Fort Edward	W17	Create a new water district to provide municipal drinking water to residents and farms along the Hudson River, from Blackhouse Road to Fort Miller.	Hudson River CARE

Table 1: Hudson River Restoration Project Proposals from the Public

County	Town or Municipality	Project #	Project Description	Submitted By
Washington	Fort Edward	W18	Repair sections of the stormwater runoff lines and construct a treatment facility to decontaminate runoff water of PCBs and TCE before the water reaches the Hudson River at the Yacht Basin.	Hudson River CARE
Washington	Hudson Falls	W19	Utilize the current General Electric office building to create a regional industrial museum.	Village of Hudson Falls
Washington	Hudson Falls	W20	Construct a park at Baker's Falls, including a viewing dock, boat launch, historical signage, and a parking area.	Village of Hudson Falls
Washington	Hudson Falls	W21	Restore and enhance Wall Street Village Pond to allow for year-round recreation. Replace vegetation, improve native habitat, and develop nature trail.	Village of Hudson Falls
Washington	Hudson Falls	W22	Supply potable municipal water (from the town of Queensbury, NY) to residents of Hudson Falls who have been denied use of local aquifer because of PCBs.	Village of Hudson Falls
Washington	Schuylerville	W23	Reimburse the Village of Schuylerville \$29,460.05 for costs incurred by removing and disposing of PCB-contaminated dredge spoil removed from boat launch site.	Village of Schuylerville
Westchester	Location not specified.	W24	Identify and remove flow and fishery impediments in Hudson River tributary Pocantico River, to improve habitat for fish and increase recreational fishing.	New York Rivers United
Location not specified.	Location not specified.	L1	Create vegetated buffers to minimize public's access to polluted areas.	Member of the Public ^b
Location not specified.	Location not specified.	L2	Enhance pollutant uptake through planting vegetation near effluent outlets on the Hudson River.	Member of the Public ^b
Location not specified.	Location not specified.	L3	Decrease bank erosion through planting vegetation in sensitive areas.	Member of the Public ^b
Location not specified.	Location not specified.	L4	Increase and improve habitat for fish through planting vegetation.	Member of the Public ^b
Location not specified.	Location not specified.	L5	Implement estuary plan.	NY DEC Hudson River Estuary Coordinator
Location not specified.	Location not specified.	L6	Restore Striped Bass fishery and spawning population.	Member of the Public ^d

Table 1: Hudson River Restoration Project Proposals from the Public

County	Town or Municipality	Project #	Project Description	Submitted By
Location not specified.	Location not specified.	L7	Dredge PCBs from river and treat to render harmless, to benefit benthic habitats.	Member of the Public ^e
Location not specified.	Location not specified.	L8	Restore wetland hydraulics to benefit habitats in tidal and nontidal wetlands.	Member of the Public ^e
Location not specified.	Location not specified.	L9	Remove obstructions and barriers to benefit fish migration.	Member of the Public ^e
Location not specified.	Location not specified.	L10	Purchase land for buffers and greenways.	Member of the Public ^e
Location not specified.	Location not specified.	L11	Create new access points and improve the public's access to the Hudson River.	Member of the Public ^e
Location not specified.	Location not specified.	L12	Purchase or otherwise acquire farmland for conservation.	Hudson River Coalition of Conservation Districts
Location not specified.	Location not specified.	L13	Partner with existing agricultural buffer program to implement urban/suburban stream buffers to protect the Hudson River.	Hudson River Coalition of Conservation Districts
Location not specified.	Location not specified.	L14	Implement a program to reduce and monitor invasive species.	Columbia County EMC
Location not specified.	Location not specified.	L15	Create more public access points for the Hudson River.	Columbia County EMC
Location not specified.	Location not specified.	L16	Work with outdoor sports associations that have ongoing restoration projects, to request conservation and education projects as well.	Columbia County EMC
Location not specified.	Location not specified.	L17	Create river access points for each community and town along the Hudson River.	Sierra Club NE Office
Location not specified.	Location not specified.	L18	Create an interpretive education center focused on conservation.	Sierra Club NE Office
Location not specified.	Location not specified.	L19	Build public recreation facilities to provide alternatives to shoreline recreation in waterfront neighborhoods.	Arbor Hill Environmental Justice Corp

Table 1: Hudson River Restoration Project Proposals from the Public

County	Town or Municipality	Project #	Project Description	Submitted By
Location not specified.	Location not specified.	L20	Create and maintain culturally sensitive educational signs at known access points along the Hudson River to provide information about fish advisories to people of color. Fund water monitoring program that engages local residents, and includes scientific experts and environmental health communications experts to translate complex information for local residents.	W.Haywood Burns Environmental Education Center
Location not specified.	Location not specified.	L21	Develop, test, and implement ecologically sound methods for restoring and managing common reed (Phragmites) stands in marshes both on and off the Hudson.	Hudsonia, Ltd.
Location not specified.	Location not specified.	L22	Construct detention basins.	Clearwater
Location not specified.	Location not specified.	L23	Limit the use of herbicides by railroad environmental engineers.	Clearwater
Location not specified.	Location not specified.	L24	Monitor the location and population of rare or endangered species communities.	Clearwater

Table 2: Hudson River Informal* Restoration Project Proposals.

Project Description	Submitted By
Address the loss of fish consumption by establishing community vegetable gardens in low income/minority neighborhoods.	Arbor Hill Environmental Justice Corp
Implement erosion control measures (such as rip-rap) to slow the migration of sediments.	Clearwater
Monitor sewage loading in the Hudson River.	Clearwater
Implement measures to control nonpoint source pollution.	Clearwater
Plant native species or otherwise establish natural vegetation within the Hudson River.	Clearwater
Repair damaged seawalls.	Clearwater
Restore both upland and tidal marshes.	Clearwater
Add baffles (barriers to flow) to create pools at low flows, which improve habitat.	Clearwater
Provide habitat elements suitable for raptors.	Clearwater
Establish program to reduce and monitor agricultural runoff.	Clearwater
Create artificial nesting structures.	Clearwater
Restore habitat for waterfowl.	Clearwater
Establish buffers in urban and suburban streams.	Clearwater
Clean up abandoned brownfields.	Clearwater
Implement a program to reduce and monitor farm runoff.	Columbia County EMC
Establish greenways to benefit riparian habitats.	Environmental Advocates
Convert waterfront brownfields in to parks.	Environmental Advocates
Secure conservation easements on farms and working forests.	Environmental Advocates
Develop and implement projects to address invasive zebra mussels and water chestnuts.	Environmental Advocates
Provide technical assistance to planning boards to evaluate the impacts of local land use projects on habitat and water quality.	Hudson River Coalition of Conservation Districts
Provide environmental education and stewardship programs for children.	Hudson River Coalition of Conservation Districts
Improve access to the Hudson River.	Scenic Hudson
Clean up abandoned brownfields to improve water quality and recreational access.	Scenic Hudson
Restore herring runs to rivers.	Scenic Hudson
Create a Hudson Riverkeeper organization for the Upper Hudson River.	Sierra Club NE Office
Implement projects to address invasive Purple loosestrife and other species.	The Nature Conservancy
Preserve habitats with significant biodiversity and/or rare and endangered species (such as Estuary Beggar-ticks and Heartleaf Plantain).	The Nature Conservancy
Preserve freshwater intertidal habitats.	The Nature Conservancy

*Informal restoration project proposals were suggested at public meetings or were suggested without a formal restoration proposal form.

FACT SHEET

HUDSON RIVER

RESTORATION PLANNING



Upper Hudson Freshwater Mussel Restoration Planning Pilot Study

Past and continuing discharges of polychlorinated biphenyls (PCBs) have contaminated Hudson River natural resources. While the U.S. Environmental Protection Agency is continuing with cleanup activities, federal and state trustee agencies – National Oceanic and Atmospheric Administration, the U.S. Department of the Interior, and New York State (the Trustees) – are assessing how releases of PCBs from the General Electric Company (GE) plants at Fort Edward and Hudson Falls, New York may have harmed the Hudson River's natural resources.

The Trustees are studying the effects of the PCB contamination through a process known as Natural Resource Damage Assessment (NRDA). In a NRDA, the Trustees use scientific studies to measure the ways hazardous substances can injure natural resources. Restoration is the goal of NRDA.

This fact sheet provides information on a freshwater mussel restoration planning pilot study being conducted as part of the Hudson River NRDA. Mussels are important components of the Hudson River ecosystem; mussels filter the water, cycle nutrients, stabilize sediments, enhance habitat complexity, and are food for wildlife.

Dredging activities in the Upper Hudson River are destroying mussel beds and mussel habitat, which are not being replaced as part of the remedy for the Hudson River PCBs Superfund Site. The proposed pilot study would collect information about mussels in areas where dredging has not yet occurred (for example: Fort Miller Pool, Stillwater Pool) starting as soon as possible, likely in late summer 2013. Other areas may be surveyed, in consultation with the Trustees, depending on dredging schedules and timing of mussel surveys. This information will serve to inform restoration planning relating to mussels adversely impacted by remedial work. Results should inform the determination of impacts of the remedy upon natural resources, spatial/temporal recovery of impacted mussel beds, and establishment of (replacement) mussel beds.

Depending on the results of this pilot study, the Trustees may conduct additional investigations focused on mussels.



Eastern Elliptio (*Elliptio complanata*);
photo credit: Phillip Westcott;
[http://digitalmedia.fws.gov/cdm/singl
eitem/collection/natdiglib/id/9733](http://digitalmedia.fws.gov/cdm/singl
eitem/collection/natdiglib/id/9733)



More Information

The following Trustee websites contain a variety of additional reports and documents relating to the overall Hudson River NRDA:

<http://www.fws.gov/contaminants/restorationplans/HudsonRiver/index.html>

<http://www.dec.ny.gov/lands/25609.html>

<http://www.darrp.noaa.gov/northeast/hudson/>

To add yourself to the **Hudson-NRDA** listserv:

1. Send a message to: requests@willamette.nos.noaa.gov
2. Write in subject: Subscribe hudsonnrda

If you have questions about natural resource damages, or want to submit a restoration project proposal, please contact one of the individuals listed below:

Tom Brosnan

National Oceanic and Atmospheric Administration

1305 East West Highway SSMC4, Room 10219

Silver Spring, MD 20910

301-713-3038 x186

Tom.Brosnan@noaa.gov

Kathryn Jahn

United States Fish and Wildlife Service

3817 Luker Road

Cortland, NY 13045

607-753-9334

Kathryn_Jahn@fws.gov

Sean Madden

New York State Department of Environmental Conservation

625 Broadway, 5th Floor

Albany, NY 12233

518-402-8977

ssmadden@gw.dec.state.ny.us

